

Anomalous Higgs Couplings in the $SO(5) \times U(1)_{B-L}$ Gauge-Higgs Unification in Warped Spacetime

Yutaka HOSOTANI^{*)} and Yutaka SAKAMURA^{**)}

Department of Physics, Osaka University, Toyonaka, Osaka 560-0043, Japan

The gauge couplings WWZ , $WWWW$, and $WWZZ$ in the gauge-Higgs unification scenario in the Randall-Sundrum warped spacetime remain almost universal as in the standard model, but substantial deviation results for the Higgs couplings. In the $SO(5) \times U(1)_{B-L}$ model, the couplings WWH and ZZH are suppressed by a factor $\cos \theta_H$ from the values in the standard model, while the bare couplings $WWHH$ and $ZZHH$ are suppressed by a factor $1 - \frac{2}{3} \sin^2 \theta_H$. Here θ_H is the Yang-Mills AB phase (Wilson line phase) along the fifth dimension, which characterizes the electroweak symmetry breaking. The suppression can be used to test the gauge-Higgs unification scenario at LHC and ILC. It is also shown that the WWZ coupling in flat spacetime deviates from the standard model value at moderate values of θ_H , contradicting with the LEP2 data.

§1. Introduction

In the previous paper we showed that substantial deviation in the Higgs couplings to W and Z bosons from those in the standard model is expected as a general feature in the gauge-Higgs unification model in warped spacetime,¹⁾ which can be tested at LHC and ILC in the coming future. Further the deviation in the WWZ coupling was shown to be very small in warped spacetime. In the present paper we give more thorough analysis of these couplings to strengthen the statements, in addition to give detailed account of the mass spectrum and wave functions in the gauge-Higgs sector in the $SO(5) \times U(1)_{B-L}$ model.

In the gauge-Higgs unification scenario the Higgs field in four dimensions is identified with the zero mode of the extra-dimensional component of gauge potentials in higher dimensional gauge theory. As such, the mass and couplings of the Higgs field are not arbitrary parameters in theory. They follow from the gauge principle. The original proposal by Fairlie and by Manton to unify the Higgs field in the six-dimensional gauge theory with S^2 as extra dimensions was unsatisfactory as it gives too low Kaluza-Klein energy scale and unrealistic couplings and spectrum.^{2), 3)} Shortly after their proposal it has been recognized that Wilson line phases, or Yang-Mills Aharonov-Bohm phases associated with non-simply connected extra dimensions can serve as the Higgs field in four dimensions. These phases, denoted as θ_H in the present paper, label classically degenerate vacua. The value of θ_H is determined at the quantum level. When the value is nontrivial, the gauge symmetry is dynamically broken.^{4), 5)}

The scenario of identifying θ_H with a 4D Higgs field, which was applied first to GUT and then to the electroweak interactions, has many attractive features.^{6)–44)}

^{*)} E-mail: hosotani@het.phys.sci.osaka-u.ac.jp

^{**)} E-mail: sakamura@het.phys.sci.osaka-u.ac.jp

Besides inducing dynamical gauge symmetry breaking, it predicts a finite mass for the Higgs field, independent of the cutoff scale.^{45)–48)} In the electroweak theory, it can solve the gauge hierarchy problem.⁷⁾ The dynamically determined value of θ_H depends on the details of the theory, particularly in the fermion sector. Astonishingly many of the features in the gauge-Higgs sector such as the mass spectrum and couplings are determined once the value of θ_H is specified. In this respect our analysis is robust. It is shown below that the WWH , ZZH , $WWHH$ and $ZZHH$ couplings are suppressed compared with those in the standard model. The predictions obtained for the gauge-Higgs couplings can be tested at LHC and ILC. If the deviation from the standard model is observed as indicated in the gauge-Higgs unification scenario, then it gives strong hint for the existence of extra dimensions. It is also confirmed that the WWZ coupling remains universal in warped spacetime, but it becomes smaller in flat spacetime compared with that in the standard model, thus already contradicting with the LEP2 data on W pair production. This strongly suggests that the extra-dimensional space is curved and warped, if it exists. There seems intimate connection between the gauge-Higgs unification scenario and the holography in the warped space.^{31), 33), 39), 42), 43)}

The paper is organized as follows. The $SO(5) \times U(1)_{B-L}$ model is set up in the next section. The spectrum and mode functions of gauge bosons are given in Sec. 3, whereas those of fermions are given in Sec. 4. Approximate masses and wave functions of gauge fields, Higgs field, and light fermions in four dimensions are given in a simple form in Sec. 5. Gauge couplings and Higgs couplings are evaluated in Sec. 6 to make predictions described above. Gauge couplings of fermions are briefly discussed in Sec. 7 and a summary is given in Sec. 8. Useful formulas are collected in appendices.

§2. $SO(5) \times U(1)_{B-L}$ model

We consider an $SO(5) \times U(1)_{B-L}$ gauge theory in the warped five-dimensional spacetime.³³⁾ The fifth dimension is compactified on an orbifold S^1/Z_2 with a radius R . We use, throughout the paper, $M, N, \dots = 0, 1, 2, 3, 4$ for the 5D curved indices, $A, B, \dots = 0, 1, 2, 3, 4$ for the 5D flat indices in tetrads, and $\mu, \nu, \dots = 0, 1, 2, 3$ for 4D flat indices. The background metric is given by⁴⁹⁾

$$ds^2 = G_{MN} dx^M dx^N = e^{-2\sigma(y)} \eta_{\mu\nu} dx^\mu dx^\nu + dy^2, \quad (2.1)$$

where $\eta_{\mu\nu} = \text{diag}(-1, 1, 1, 1)$, $\sigma(y) = \sigma(y + 2\pi R)$, and $\sigma(y) \equiv k|y|$ for $|y| \leq \pi R$. The cosmological constant in the bulk 5D spacetime is given by $\Lambda = -k^2$. $(x^\mu, -y)$ and $(x^\mu, y + 2\pi R)$ are identified with (x^μ, y) . The spacetime is equivalent to the interval in the fifth dimension y with two boundaries at $y = 0$ and $y = \pi R$, which we refer to as the Planck brane and the TeV brane, respectively.

There are $SO(5)$ gauge fields A_M and $U(1)_{B-L}$ gauge field B_M . The former are decomposed as

$$A_M = \sum_{I=1}^{10} A_M^I T^I = \sum_{a_L=1}^3 A_M^{a_L} T^{a_L} + \sum_{a_R=1}^3 A_M^{a_R} T^{a_R} + \sum_{\hat{a}=1}^4 A_M^{\hat{a}} T^{\hat{a}}, \quad (2.2)$$

where T^{a_L, a_R} ($a_L, a_R = 1, 2, 3$) and $T^{\hat{a}}$ ($\hat{a} = 1, 2, 3, 4$) are the generators of $SO(4) \sim SU(2)_L \times SU(2)_R$ and $SO(5)/SO(4)$, respectively. The spinorial representation of T^I is tabulated in (A.1) in appendix A. As a matter field we introduce a spinor field Ψ in the spinorial representation of $SO(5)$ (*i.e.*, **4** of $SO(5)$) in the bulk as an example.

The relevant part of the action in the bulk is^{50),51)}

$$S = \int d^5x \sqrt{-G} \left[-\text{tr} \left(\frac{1}{2} F^{(A)MN} F_{MN}^{(A)} + \frac{1}{\xi} (f_{\text{gf}}^{(A)})^2 + \mathcal{L}_{\text{gh}}^{(A)} \right) \right. \\ \left. - \left(\frac{1}{4} F^{(B)MN} F_{MN}^{(B)} + \frac{1}{2\xi} (f_{\text{gf}}^{(B)})^2 + \mathcal{L}_{\text{gh}}^{(B)} \right) + i\bar{\Psi} \Gamma^N \mathcal{D}_N \Psi - iM_\Psi \varepsilon \bar{\Psi} \Psi \right], \quad (2.3)$$

where $G \equiv \det(G_{MN})$ and $\Gamma^N \equiv e_A^N \Gamma^A$. The 5D γ -matrices Γ^A are related to the 4D ones γ^μ by $\Gamma^\mu = \gamma^\mu$ and $\Gamma^4 = \gamma_5$ which is the 4D chiral operator. The gauge-fixing functions $f_{\text{gf}}^{(A,B)}$ are specified in the next section. $\mathcal{L}_{\text{gh}}^{(A,B)}$ are the associated ghost Lagrangians, and M_Ψ is a bulk (kink) mass parameter. Since the operator $\bar{\Psi}\Psi$ is Z_2 -odd, we need the periodic sign function $\varepsilon(y) = \sigma'(y)/k$ satisfying $\varepsilon(y) = \pm 1$. The field strengths and the covariant derivatives are defined by

$$F_{MN}^{(A)} \equiv \partial_M A_N - \partial_N A_M - ig_A [A_M, A_N], \\ F_{MN}^{(B)} \equiv \partial_M B_N - \partial_N B_M, \\ \mathcal{D}_M \Psi \equiv \left\{ \partial_M - \frac{1}{4} \omega_M^{AB} \Gamma_{AB} - ig_A A_M - i\frac{g_B}{2} \mathcal{Q}_{\text{B-L}} B_M \right\} \Psi, \quad (2.4)$$

where g_A (g_B) is the 5D gauge coupling for A_M (B_M), $\mathcal{Q}_{\text{B-L}}$ is a charge of $U(1)_{\text{B-L}}$, and $\Gamma^{AB} \equiv \frac{1}{2} [\Gamma^A, \Gamma^B]$. The spin connection 1-form $\omega_M^{AB} = \omega_M^{AB} dx^M$ determined from the metric (2.1) is

$$\omega^{\nu 4} = -\sigma' e^{-\sigma} dx^\nu, \quad \text{other components} = 0. \quad (2.5)$$

The boundary conditions at the fixed points $y_0 = 0$ and $y_\pi = \pi R$, which preserve the orbifold structure, are

$$\begin{pmatrix} A_\mu \\ A_y \end{pmatrix} (x, y_j - y) = P_j \begin{pmatrix} A_\mu \\ -A_y \end{pmatrix} (x, y_j + y) P_j^{-1}, \\ \begin{pmatrix} B_\mu \\ B_y \end{pmatrix} (x, y_j - y) = \begin{pmatrix} B_\mu \\ -B_y \end{pmatrix} (x, y_j + y), \\ \Psi(x, y_j - y) = \eta_j P_j \gamma_5 \Psi(x, y_j + y), \quad (2.6)$$

where $\eta_j = \pm 1$ ($j = 0, \pi$). $P_j \in SO(5)$ are constant matrices satisfying $P_j^2 = 1$. In the present paper we take

$$P_0 = P_\pi = \begin{pmatrix} 1_2 & \\ & -1_2 \end{pmatrix} \quad (2.7)$$

in the spinorial representation, or equivalently $P_0 = P_\pi = \text{diag}(-1, -1, -1, -1, 1)$ in the vectorial representation. Then the gauge symmetry is broken to $SO(4) \times U(1)_{\text{B-L}} \sim SU(2)_L \times SU(2)_R \times U(1)_{\text{B-L}}$ at both boundaries. (The broken generators are $T^{\hat{a}}$; $\hat{a} = 1, 2, 3, 4$.) It is convenient to decompose Ψ as

$$\Psi = \begin{pmatrix} q \\ Q \end{pmatrix}, \quad (2.8)$$

where q and Q belong to $(\frac{1}{2}, 0)$ and $(0, \frac{1}{2})$ of $SU(2)_L \times SU(2)_R$, respectively.

Fields with Neumann boundary conditions at both boundaries have zero modes when perturbation theory is developed around the trivial configuration $A_M = 0$. With (2.6) and (2.7) there arise zero modes for $A_y^{\hat{a}}$ ($\hat{a} = 1, 2, 3, 4$). They are identified with the $SU(2)_L$ doublet-Higgs field in the standard model; $\Phi \propto (A_y^{\hat{1}} + iA_y^{\hat{2}}, A_y^{\hat{4}} - iA_y^{\hat{3}})^t$. A nonvanishing expectation value of $A_y^{\hat{4}}$ gives rise to a Wilson line phase or Yang-Mills Aharonov-Bohm phase, $\theta_H \equiv g_A \int_0^{2\pi R} dy \langle A_y^{\hat{4}} \rangle / 2\sqrt{2} = (g_A/\sqrt{2}) \int_0^{\pi R} dy \langle A_y^{\hat{4}} \rangle$. More explicitly,

$$\langle A_y^{\hat{4}} \rangle = \frac{2\sqrt{2}ke^{2ky}}{g_A(e^{2k\pi R} - 1)} \theta_H. \quad (2.9)$$

Although $\theta_H \neq 0$ gives vanishing field strengths, it affects physics at the quantum level. The global minimum of the effective potential for θ_H determines the quantum vacuum.⁴⁾ The nonvanishing θ_H induces dynamical electroweak gauge symmetry breaking.

There are residual gauge transformations which maintain the boundary condition (2.6).^{5), 36)} A large gauge transformation given by

$$\Omega^{\text{large}}(y) = \exp \left\{ in\pi \frac{e^{2ky} - 1}{e^{2k\pi R} - 1} (2\sqrt{2} T^{\hat{4}}) \right\} \quad (2.10)$$

($0 \leq y \leq \pi R$, n : an integer) shifts θ_H by $\theta_H + 2\pi n$, which implies that all physical quantities are periodic functions of θ_H . The large gauge invariance is vital to guarantee the finiteness of the Higgs boson mass.^{45), 48)} The θ_H -dependent part of the effective potential diverges without the large gauge invariance.

The even-odd property in (2.6) does not completely fix boundary conditions of the fields. If there are no additional dynamics on the two branes, fields which are odd under parity at $y = 0$ or πR obey the Dirichlet boundary condition (D) so that they vanish there. On the other hand, fields which are even under parity obey the Neumann boundary conditions (N). For gauge fields the Neumann boundary condition is given by $dA_\mu/dy = 0$ or $d(e^{-2ky}A_y)/dy = 0$. As a result of additional dynamics on the branes, however, a field with even parity, for instance, can effectively obey the Dirichlet boundary condition. The field develops a cusp-type singularity there due to brane dynamics. As discussed below, the $SO(4)$ symmetry on the Planck brane is broken to $SU(2)_L \times U(1)_Y$ in this manner.

Let us define new fields A_M^{3R} and A_M^Y by

$$\begin{pmatrix} A_M^{3R} \\ A_M^Y \end{pmatrix} = \begin{pmatrix} c_\phi & -s_\phi \\ s_\phi & c_\phi \end{pmatrix} \begin{pmatrix} A_M^{3R} \\ B_M \end{pmatrix},$$

Table I. Boundary conditions for the gauge fields. $a_L, a_R = 1, 2, 3$ and $\hat{a} = 1, 2, 3, 4$. The notation (D,N), for example, denotes the Dirichlet boundary condition at $y = 0$ and the Neumann boundary condition at $y = \pi R$.

$A_\mu^{a_L}$	$A_\mu^{1,2R}$	$A_\mu'^{3R}$	A_μ^Y	$A_\mu^{\hat{a}}$	$A_y^{a_L}$	$A_y^{1,2R}$	$A_y'^{3R}$	A_y^Y	$A_y^{\hat{a}}$
(N,N)	(D,N)	(D,N)	(N,N)	(D,D)	(D,D)	(D,D)	(D,D)	(D,D)	(N,N)

$$c_\phi \equiv \frac{g_A}{\sqrt{g_A^2 + g_B^2}} \quad , \quad s_\phi \equiv \frac{g_B}{\sqrt{g_A^2 + g_B^2}} \quad . \quad (2.11)$$

$A_\mu^{a_R}$ and B_μ are even under parity, whereas $A_y^{a_R}$ and B_y are odd. We suppose that as a result of additional dynamics on the Planck brane the even fields A_μ^{1R} , A_μ^{2R} , and $A_\mu'^{3R}$ obey the Dirichlet (D) boundary condition there. The boundary conditions for gauge fields are tabulated in Table I. It is confirmed that the boundary conditions in Table I preserve the large gauge invariance, that is, new gauge potentials obtained by (2.10) obey the same boundary conditions as the original fields. We remark that the Neumann (N) boundary condition on the Planck brane cannot be imposed on A_y^{1R} , A_y^{2R} , and $A_y'^{3R}$, as it does not preserve the large gauge invariance. Indeed, under a large gauge transformation (2.10), $A_y \rightarrow A_y'$ and

$$\frac{d}{dy} \left(e^{-2ky} A_y^{\hat{a}} \right) = \frac{d}{dy} \left(e^{-2ky} A_y^{\hat{a}} \right) + \frac{2\sqrt{2} n \pi k}{e^{2\pi k R} - 1} (A_y^{a_L} - A_y^{a_R}) \quad (2.12)$$

for $a = 1, 2, 3$ so that the Neumann boundary condition for $A_y^{\hat{a}}$ ($a = 1, 2, 3$) is preserved only if $A_y^{a_L} = A_y^{a_R}$ at the boundary. Since $A_y^{a_L}$ obeys the Dirichlet boundary condition, $A_y^{a_R}$ must obey the Dirichlet boundary condition as well.

With the boundary condition in Table I, the gauge symmetry $SO(5) \times U(1)_{B-L}$ in the bulk is reduced to $SO(4) \times U(1)_{B-L}$ at the TeV brane and to $SU(2)_L \times U(1)_Y$ at the Planck brane. The resultant symmetry of the theory is $SU(2)_L \times U(1)_Y$, which is subsequently broken to $U(1)_{EM}$ by nonvanishing $A_y^{\hat{a}}$ ($\hat{a} = 1, 2, 3, 4$) or θ_H . The weak hypercharge Y is given by $Y = T^{3R} + q_{B-L}/2$.

The boundary conditions of A_μ^{1R} , A_μ^{2R} , and $A_\mu'^{3R}$ are changed from N to D on the Planck brane, if additional dynamics on the Planck brane spontaneously breaks $SU(2)_R \times U(1)_{B-L}$ to $U(1)_Y$ at relatively high energy scale M , say, near the Planck scale M_{Pl} so that A_μ^{1R} , A_μ^{2R} , and $A_\mu'^{3R}$ have masses of $O(M)$ on the Planck brane. Below the TeV scale, the mass terms on the Planck brane strongly suppress the boundary values of these fields, effectively changing the boundary conditions from N to D at the Planck brane. With the underlying gauge invariance it is expected that the tree unitarity for the gauge boson scatterings^{53)–55)} is preserved with these boundary conditions.

§3. Spectrum and mode functions of gauge bosons

In this section we derive the spectrum and mode functions of gauge fields with the boundary conditions listed in Table I. Although such quantities have been well

discussed in many papers in the case of vanishing θ_H (see Refs. 50)–52) for example), the case of nonzero θ_H becomes highly nontrivial as the boundary conditions of 5D fields are twisted by the angle θ_H , *i.e.*, they are no longer either the ordinary Neumann or Dirichlet boundary conditions. In fact the $SU(3)$ model has been analysed in Ref. 37) where it is found that the wave functions have nontrivial θ_H dependence. Here and in the next section we provide systematic KK analysis and obtained the full spectrum and the wave functions in the $SO(5) \times U(1)_{B-L}$ model for a general value of θ_H . The results obtained here are used to estimate various coupling constants in Sec. 6.

3.1. General solutions in the bulk

The basic procedure is the same as in our previous papers.^{1),37)} We employ the background field method, separating A_M (B_M) into the classical part A_M^c (B_M^c) and the quantum part A_M^q (B_M^q); $A_M = A_M^c + A_M^q$ and $B_M = B_M^c + B_M^q$. It is convenient to go over to the conformal coordinate $z \equiv e^{\sigma(y)}$ for the fifth dimension;

$$ds^2 = \frac{1}{z^2} \left\{ \eta_{\mu\nu} dx^\mu dx^\nu + \frac{dz^2}{k^2} \right\},$$

$$\partial_y = kz \partial_z, \quad A_y = kz A_z, \quad B_y = kz B_z. \quad (3.1)$$

The boundaries are located at $z = 1$ and $z_\pi \equiv e^{k\pi R}$. The gauge-fixing functions are chosen³⁵⁾ as

$$f_{\text{gf}}^{(A)} = z^2 \left\{ \eta^{\mu\nu} \mathcal{D}_\mu^c A_\nu^q + \xi k^2 z \mathcal{D}_z^c \left(\frac{1}{z} A_z^q \right) \right\},$$

$$f_{\text{gf}}^{(B)} = z^2 \left\{ \eta^{\mu\nu} \partial_\mu B_\nu^q + \xi k^2 z \partial_z \left(\frac{1}{z} B_z^q \right) \right\} \quad (3.2)$$

where $\mathcal{D}_M^c A_N^q \equiv \partial_M A_N^q - ig_A [A_M^c, A_N^q]$.

The quadratic terms for the gauge and ghost fields are simplified for $\xi = 1$,

$$S = \int d^4x \frac{dz}{kz} \left[\text{tr} \left\{ \eta^{\mu\nu} A_\mu^q (\square + k^2 \mathcal{P}_4) A_\nu^q + k^2 A_z^q (\square + k^2 \mathcal{P}_z) A_z^q + \mathcal{L}_{\text{gh}}^{(A)} \right\} \right. \\ \left. + \eta^{\mu\nu} B_\mu^q (\square + k^2 \mathcal{P}_4) B_\nu^q + k^2 B_z^q (\square + k^2 \mathcal{P}_z) B_z^q + \mathcal{L}_{\text{gh}}^{(B)} \right], \quad (3.3)$$

where

$$\square \equiv \eta^{\mu\nu} \partial_\mu \partial_\nu, \quad \mathcal{P}_4 \equiv z \mathcal{D}_z^c \frac{1}{z} \mathcal{D}_z^c, \quad \mathcal{P}_z \equiv \mathcal{D}_z^c z \mathcal{D}_z^c \frac{1}{z}. \quad (3.4)$$

Here we have taken $A_\mu^c = 0$, respecting the 4D Poincaré symmetry. The surface terms at the boundaries at $z = 1, z_\pi$ vanish thanks to the boundary conditions for each field.

The linearized equations of motion for A_M become

$$\square A_\mu^q + k^2 z \mathcal{D}_z^c \frac{1}{z} \mathcal{D}_z^c A_\mu^q = 0,$$

$$\square A_z^q + k^2 \mathcal{D}_z^c z \mathcal{D}_z^c \frac{1}{z} A_z^q = 0, \quad (3.5)$$

and those for B_M have similar forms. The classical background is taken to be $A_z^c = vzT^{\hat{4}}$ (v : real constant) below.

We move to a twisted basis by a gauge transformation;^{*)}

$$\begin{aligned}\tilde{A}_M &\equiv \Omega A_M^q \Omega^{-1}, \quad \tilde{B}_M \equiv B_M^q, \\ \Omega(z) &\equiv \exp \left\{ ig_A \int_z^{z_\pi} dz' A_z^c(z') \right\}.\end{aligned}\quad (3.6)$$

As shown in Refs. 5), 56), 57), sets of boundary conditions related by gauge transformations form equivalence classes and all sets in each equivalence class are physically equivalent. Dynamics of the Yang-Mills AB phase θ_H guarantees the equivalence. In the twisted basis the classical background of the gauge fields vanishes so that the linearized equations of motion reduce to free field equations, while the boundary conditions become more involved.

$$\begin{aligned}\square \tilde{A}_\mu + k^2 \left(\partial_z^2 - \frac{1}{z} \partial_z \right) \tilde{A}_\mu &= 0 \quad , \quad \square \tilde{A}_z + k^2 \left(\partial_z^2 - \frac{1}{z} \partial_z + \frac{1}{z^2} \right) \tilde{A}_z = 0 \quad , \\ \square \tilde{B}_\mu + k^2 \left(\partial_z^2 - \frac{1}{z} \partial_z \right) \tilde{B}_\mu &= 0 \quad , \quad \square \tilde{B}_z + k^2 \left(\partial_z^2 - \frac{1}{z} \partial_z + \frac{1}{z^2} \right) \tilde{B}_z = 0 \quad .\end{aligned}\quad (3.7)$$

Thus the equations for eigenmodes with a mass eigenvalue $m_n = k\lambda_n$ are

$$\begin{aligned}\left\{ \frac{d^2}{dz^2} - \frac{1}{z} \frac{d}{dz} + \lambda_n^2 \right\} \tilde{h}_{A,n}^I &= 0 \quad , \\ \left\{ \frac{d^2}{dz^2} - \frac{1}{z} \frac{d}{dz} + \frac{1}{z^2} + \lambda_n^2 \right\} \tilde{h}_{\varphi,n}^I &= 0 \quad .\end{aligned}\quad (3.8)$$

With these eigenfunctions the gauge potentials are expanded as

$$\begin{aligned}\tilde{A}_\mu^I(x, z) &= \sum_n \tilde{h}_{A,n}^I(z) A_\mu^{(n)}(x) \quad , \quad \tilde{A}_z^I(x, z) = \sum_n \tilde{h}_{\varphi,n}^I(z) \varphi^{(n)}(x) \quad , \\ \tilde{B}_\mu(x, z) &= \sum_n \tilde{h}_{A,n}^B(z) A_\mu^{(n)}(x) \quad , \quad \tilde{B}_z(x, z) = \sum_n \tilde{h}_{\varphi,n}^B(z) \varphi^{(n)}(x) \quad .\end{aligned}\quad (3.9)$$

The general solutions to Eq.(3.8) are expressed in terms of the Bessel functions as

$$\begin{aligned}\tilde{h}_{A,n}^{\bar{I}}(z) &= z \left\{ \alpha_{A,n}^{\bar{I}} J_1(\lambda_n z) + \beta_{A,n}^{\bar{I}} Y_1(\lambda_n z) \right\} \quad , \\ \tilde{h}_{\varphi,n}^{\bar{I}}(z) &= z \left\{ \alpha_{\varphi,n}^{\bar{I}} J_0(\lambda_n z) + \beta_{\varphi,n}^{\bar{I}} Y_0(\lambda_n z) \right\} \quad ,\end{aligned}\quad (3.10)$$

where $\bar{I} = (a_L, a_R, \hat{a}, B)$ and α_n 's and β_n 's are constants to be determined by the boundary conditions.

^{*)} We define $\Omega(z)$ so that $\Omega(z_\pi) = 1_4$ in contrast to our previous work^{1), 37)} where it is defined as $\Omega(1) = 1_4$.

3.2. Mass eigenvalues and mode functions

To determine the eigenvalues λ_n 's and the corresponding mode functions (3·10), we need to take into account the boundary conditions listed in Table I. In this subsection we mainly examine the 4D components of the gauge fields (A_μ , B_μ). The mass spectrum and the mode functions for the extra-dimensional components (A_z, B_z) are examined in the next subsection.

At the boundaries

$$\partial_z A_\mu^{aL} = \partial_z A_\mu^Y = 0 \quad , \quad A_\mu^{1,2R} = A_\mu^{3R} = A_\mu^{\hat{a}} = 0 \quad \text{at } z = 1 \quad , \quad (3\cdot11)$$

$$\partial_z A_\mu^{aL} = \partial_z A_\mu^{aR} = \partial_z B_\mu = 0 \quad , \quad A_\mu^{\hat{a}} = 0 \quad \text{at } z = z_\pi \quad . \quad (3\cdot12)$$

We translate these conditions into those in the twisted basis ($\tilde{A}_M, \tilde{B}_M$). Among the extra-dimensional components of the gauge fields, only $A_z^{\hat{a}}$ can have non-vanishing vacuum expectation values (VEV). With the residual $SU(2)_L \times U(1)_Y$ symmetry, we can restrict ourselves to

$$A_z^c = v z T^{\hat{4}} \quad , \quad (3\cdot13)$$

where a constant v is related to θ_H by

$$\theta_H = \frac{gA v}{2\sqrt{2}}(z_\pi^2 - 1) \quad . \quad (3\cdot14)$$

The potential has a classical flat direction along θ_H . The value for θ_H is determined at the quantum level. Using (3·13), the gauge transformation matrix Ω defined in (3·6) is calculated as

$$\Omega(z) = \exp \left\{ i \frac{1}{2} \theta(z) (2\sqrt{2} T^{\hat{4}}) \right\} = \begin{pmatrix} c_\theta 1_2 & i s_\theta 1_2 \\ i s_\theta 1_2 & c_\theta 1_2 \end{pmatrix} \quad , \quad (3\cdot15)$$

where

$$\theta(z) \equiv \frac{gA}{\sqrt{2}} \int_z^{z_\pi} dz' A_z^{\hat{4}}(z') = \frac{gA v}{2\sqrt{2}}(z_\pi^2 - z^2) = \theta_H \frac{z_\pi^2 - z^2}{z_\pi^2 - 1} \quad , \quad (3\cdot16)$$

and $c_\theta \equiv \cos \frac{1}{2} \theta(z)$, $s_\theta \equiv \sin \frac{1}{2} \theta(z)$. Thus the relation between A_M and $\tilde{A}_M$ in Eq.(3·6) can be written as

$$\begin{pmatrix} \tilde{A}_M^{aL} \\ \tilde{A}_M^{aR} \\ \tilde{A}_M^{\hat{a}} \end{pmatrix} = \begin{pmatrix} c_\theta^2 & s_\theta^2 & \sqrt{2} s_\theta c_\theta \\ s_\theta^2 & c_\theta^2 & -\sqrt{2} s_\theta c_\theta \\ -\sqrt{2} s_\theta c_\theta & \sqrt{2} s_\theta c_\theta & c_\theta^2 - s_\theta^2 \end{pmatrix} \begin{pmatrix} A_M^{qaL} \\ A_M^{qaR} \\ A_M^{q\hat{a}} \end{pmatrix} \quad , \quad (3\cdot17)$$

$$\tilde{A}_M^{\hat{4}} = A_M^{q\hat{4}} \quad ,$$

where $a = 1, 2, 3$.

The boundary conditions (3·11) and (3·12) can be rewritten in terms of $\tilde{A}_\mu$ by using this relation. The condition (3·12) determines the ratios between $\alpha_{A,n}$'s and $\beta_{A,n}$'s in Eq.(3·10) so that the mode functions have the following forms.

$$\tilde{h}_{A,n}^{aL}(z) = C_{A,n}^{aL} z F_{1,0}(\lambda_n z, \lambda_n z_\pi) \quad ,$$

$$\begin{aligned}
 \tilde{h}_{A,n}^{aR}(z) &= C_{A,n}^{aR} z F_{1,0}(\lambda_n z, \lambda_n z_\pi) , \\
 \tilde{h}_{A,n}^{\hat{a}}(z) &= C_{A,n}^{\hat{a}} z F_{1,1}(\lambda_n z, \lambda_n z_\pi) , \\
 \tilde{h}_{A,n}^B(z) &= C_{A,n}^B z F_{1,0}(\lambda_n z, \lambda_n z_\pi) .
 \end{aligned} \tag{3.18}$$

Here the functions $F_{\alpha,\beta}(u, v)$ are defined in (C.1). The mass eigenvalue λ_n and the coefficients C_n 's are determined by the remaining boundary condition (3.11), which amount to

$$\begin{pmatrix} c_H^2 \lambda_n F_{0,0} & s_H^2 \lambda_n F_{0,0} & -\sqrt{2} s_H c_H F_{0,1} \\ s_H^2 \lambda_n F_{1,0} & c_H^2 \lambda_n F_{1,0} & \sqrt{2} s_H c_H F_{1,1} \\ \sqrt{2} s_H c_H \lambda_n F_{1,0} & -\sqrt{2} s_H c_H \lambda_n F_{1,0} & (c_H^2 - s_H^2) F_{1,1} \end{pmatrix} \begin{pmatrix} C_{A,n}^{aL} \\ C_{A,n}^{aR} \\ C_{A,n}^{\hat{a}} \end{pmatrix} = 0 , \tag{3.19}$$

for $a = 1, 2$, and

$$\begin{pmatrix} c_H^2 \lambda_n F_{0,0} & s_H^2 \lambda_n F_{0,0} & -\sqrt{2} s_H c_H F_{0,1} & 0 \\ c_\phi s_H^2 \lambda_n F_{1,0} & c_\phi c_H^2 \lambda_n F_{1,0} & \sqrt{2} c_\phi s_H c_H F_{1,1} & -s_\phi \lambda_n F_{1,0} \\ \sqrt{2} s_H c_H \lambda_n F_{1,0} & -\sqrt{2} s_H c_H \lambda_n F_{1,0} & (c_H^2 - s_H^2) F_{1,1} & 0 \\ s_\phi s_H^2 \lambda_n F_{0,0} & s_\phi c_H^2 \lambda_n F_{0,0} & \sqrt{2} s_\phi s_H c_H F_{0,1} & c_\phi \lambda_n F_{0,0} \end{pmatrix} \begin{pmatrix} C_{A,n}^{3L} \\ C_{A,n}^{3R} \\ C_{A,n}^3 \\ C_{A,n}^B \end{pmatrix} = 0 , \tag{3.20}$$

$$F_{1,1} C_{A,n}^4 = 0 , \tag{3.21}$$

where $s_H \equiv \sin \frac{1}{2} \theta_H$, $c_H \equiv \cos \frac{1}{2} \theta_H$. Here and henceforth, $F_{\alpha,\beta}$ without the argument denotes $F_{\alpha,\beta}(\lambda_n, \lambda_n z_\pi)$.

$U(1)$ subgroup remains unbroken for any value of nonzero θ_H , which is identified with the electromagnetic gauge group $U(1)_{\text{EM}}$. The gauge fields are classified in three sectors, the charged sector

$$(A_M^{\pm L}, A_M^{\pm R}, A_M^{\hat{\pm}}) \equiv \frac{1}{\sqrt{2}} (A_M^{1L} \pm i A_M^{2L}, A_M^{1R} \pm i A_M^{2R}, A_M^{\hat{1}} \pm i A_M^{\hat{2}}) , \tag{3.22}$$

the neutral sector

$$(A_M^{3L}, A_M^{3R}, A_M^{\hat{3}}, B_M) , \tag{3.23}$$

and the ‘singlet’ sector A_M^4 . The latter two sectors are neutral under $U(1)_{\text{EM}}$. The orthonormal relations among the mode functions are

$$\begin{aligned}
 \int_1^{z_\pi} \frac{dz}{kz} \left\{ \tilde{h}_{A,n}^{\pm L} \tilde{h}_{A,l}^{\pm L} + \tilde{h}_{A,n}^{\pm R} \tilde{h}_{A,l}^{\pm R} + \tilde{h}_{A,n}^{\hat{\pm}} \tilde{h}_{A,l}^{\hat{\pm}} \right\} &= \delta_{n,l} , \\
 \int_1^{z_\pi} \frac{dz}{kz} \left\{ \tilde{h}_{A,n}^{3L} \tilde{h}_{A,l}^{3L} + \tilde{h}_{A,n}^{3R} \tilde{h}_{A,l}^{3R} + \tilde{h}_{A,n}^{\hat{3}} \tilde{h}_{A,l}^{\hat{3}} + \tilde{h}_{A,n}^B \tilde{h}_{A,l}^B \right\} &= \delta_{n,l} , \\
 \int_1^{z_\pi} \frac{dz}{kz} \tilde{h}_{A,n}^4 \tilde{h}_{A,l}^4 &= \delta_{n,l} .
 \end{aligned} \tag{3.24}$$

3.2.1. Charged sector $(A_\mu^{\pm L}, A_\mu^{\pm R}, A_\mu^{\hat{\pm}})$

In order for nontrivial solutions to Eq. (3.19) to exist, the determinant of the 3×3 matrix must vanish, which leads to

$$F_{1,0} \left\{ \pi^2 \lambda_n^2 z_\pi F_{0,0} F_{1,1} - 2 \sin^2 \theta_H \right\} = 0 . \tag{3.25}$$

Once the spectrum λ_n is determined by the above equation, the corresponding $C_{A,n}$'s are fixed by (3.19) with the normalization condition (3.24).

There are two cases for the mass spectrum.

Case 1: $F_{1,0} = 0$

There is no massless mode and the lightest mode has a mass of $\mathcal{O}(m_{\text{KK}})$, where the Kaluza-Klein (KK) mass scale m_{KK} is given by

$$m_{\text{KK}} \equiv \frac{k\pi}{z_\pi - 1} . \quad (3.26)$$

The coefficients $C_{A,n}^I$ ($I = \pm_L, \pm_R, \pm$) in the mode functions are given by

$$\begin{aligned} C_{A,n}^{\pm_L} &= (1 - \cos \theta_H) \hat{C}_1 , \\ C_{A,n}^{\pm_R} &= -(1 + \cos \theta_H) \hat{C}_1 , \\ C_{A,n}^\pm &= 0 , \\ \hat{C}_1 &= \frac{\sqrt{k}}{\sqrt{1 + \cos^2 \theta_H}} \left\{ \frac{4}{\pi^2 \lambda_n^2} - F_{0,0}^2 \right\}^{-1/2} . \end{aligned} \quad (3.27)$$

The mass spectrum is independent of θ_H and is the same as for the modes with the boundary condition (D,N) at $\theta_H = 0$. For nonzero θ_H , however, the above modes do not have definite Z_2 -parities since components with different boundary conditions mix with each other. This can be seen explicitly from the fact that the mode functions have nontrivial θ_H -dependences.

Case 2: $\pi^2 \lambda_n^2 z_\pi F_{0,0} F_{1,1} = 2 \sin^2 \theta_H$

In this case the mass spectrum depends on θ_H . The lightest mode is massless at $\theta_H = 0$, while it acquires a nonvanishing mass when $\theta_H \neq 0$. The lightest mode is identified with the W boson. The coefficients in the mode functions are given by

$$\begin{aligned} C_{A,n}^{\pm_L} &= (1 + \cos \theta_H) \hat{C}_2 , \\ C_{A,n}^{\pm_R} &= (1 - \cos \theta_H) \hat{C}_2 , \\ C_{A,n}^\pm &= -\sqrt{2} \sin \theta_H \frac{F_{1,0}}{F_{1,1}} \hat{C}_2 , \\ \hat{C}_2 &= \frac{\sqrt{k}}{\sqrt{1 + \cos^2 \theta_H}} \left\{ \frac{4}{\pi^2 \lambda_n^2} + \frac{\pi^2 \lambda_n^2 z_\pi^2 F_{1,0}^2 F_{0,0}^2}{\sin^2 \theta_H (1 + \cos^2 \theta_H)} - \frac{2F_{1,0}^2}{1 + \cos^2 \theta_H} - \frac{2F_{0,0}^2}{\sin^2 \theta_H} \right\}^{-1/2} \end{aligned} \quad (3.28)$$

One comment is in order about the behavior of $\hat{C}_2$ in the $\theta_H \rightarrow 0$ limit. For the KK excited states, or modes with $\lim_{\theta_H \rightarrow 0} \lambda_n \neq 0$, either $F_{0,0}$ or $F_{1,1}$ is $\mathcal{O}(\theta_H^2)$. $C_{A,n}^{\pm_L}$ becomes dominant for the modes with $F_{0,0} = \mathcal{O}(\theta_H^2)$, while $C_{A,n}^\pm$ becomes dominant for the modes with $F_{1,1} = \mathcal{O}(\theta_H^2)$. For the zero mode $\lambda_0 = \mathcal{O}(\theta_H)$ and $\hat{C}_2 \simeq \sqrt{\pi} \lambda_0 / 4\sqrt{R}$.

3.2.2. Neutral sector ($A_\mu^{3L}, A_\mu^{3R}, A_\mu^{\hat{3}}, B_\mu$)

The determinant of the 4×4 matrix in (3·20) must vanish, which leads to

$$\lambda_n F_{0,0} F_{1,0} \left\{ \pi^2 \lambda_n^2 z_\pi F_{0,0} F_{1,1} - 2(1 + s_\phi^2) \sin^2 \theta_H \right\} = 0 \quad . \quad (3·29)$$

s_ϕ is defined in (2·11). This determines the mass spectrum. Once λ_n is determined, the coefficients $C_{A,n}^{\bar{I}}$ ($\bar{I} = 3_L, 3_R, \hat{3}, B$) in mode functions are fixed by (3·20) with the normalization condition (3·24).

The neutral sector is classified into three cases.

Case 1: $\lambda_n F_{0,0} = 0$

The massless mode ($\lambda_0 = 0$) identified with the photon for the unbroken $U(1)_{EM}$ has a constant mode function

$$\begin{aligned} \tilde{h}_{A,0}^{3L} &= \tilde{h}_{A,0}^{3R} = \frac{s_\phi}{\sqrt{(1 + s_\phi^2)\pi R}} \quad , \\ \tilde{h}_{A,0}^{\hat{3}} &= 0 \quad , \quad \tilde{h}_{A,0}^B = \frac{c_\phi}{\sqrt{(1 + s_\phi^2)\pi R}} \quad . \end{aligned} \quad (3·30)$$

The massive modes have

$$\begin{aligned} \tilde{h}_{A,n}^{3L}(z) &= \tilde{h}_{A,n}^{3R}(z) = \frac{\sqrt{2k}s_\phi}{\sqrt{1 + s_\phi^2}} \left\{ \frac{4}{\pi^2 \lambda_n^2} - F_{1,0}^2 \right\}^{-1/2} F_{1,0}(\lambda_n z, \lambda_n z_\pi) \quad , \\ \tilde{h}_{A,n}^{\hat{3}}(z) &= 0 \quad , \quad \tilde{h}_{A,n}^B(z) = \frac{c_\phi}{s_\phi} \tilde{h}_{A,n}^{3L}(z) \quad . \end{aligned} \quad (3·31)$$

(3·30) can be obtained also from (3·31) by taking the limit of $\lambda_n \rightarrow 0$. Note that the mass spectrum and the mode functions in the photon sector are independent of θ_H . In fact we can extract the photon sector from the neutral sector by the following field redefinition.

$$\begin{pmatrix} A_\mu^{3V} \\ A_\mu^{3A} \\ A_\mu^{\hat{3}} \\ A_\mu^\gamma \end{pmatrix} = \frac{1}{\sqrt{2(1 + s_\phi^2)}} \begin{pmatrix} \frac{c_\phi}{\sqrt{1 + s_\phi^2}} & -\frac{c_\phi}{\sqrt{1 + s_\phi^2}} & -2s_\phi \\ \sqrt{2}s_\phi & \sqrt{2}s_\phi & 0 \\ \sqrt{2}s_\phi & \sqrt{2}s_\phi & \sqrt{2}c_\phi \end{pmatrix} \begin{pmatrix} A_\mu^{3L} \\ A_\mu^{3R} \\ B_\mu \end{pmatrix} \quad . \quad (3·32)$$

The photon field A_μ^γ does not mix with the other components (A_μ^{3V}, A_μ^{3A}) under the Ω -rotation in (3·15).

Case 2: $F_{1,0} = 0$

The equation that determines the mass spectrum is the same as in the case 1 in the charged sector. The coefficients in the mode functions are given by

$$\begin{aligned} C_{A,n}^{3L} &= (1 - \cos \theta_H) \hat{C}_3 \quad , \\ C_{A,n}^{3R} &= -(1 + \cos \theta_H) \hat{C}_3 \quad , \\ C_{A,n}^{\hat{3}} &= 0 \quad , \\ C_{A,n}^B &= 2t_\phi \cos \theta_H \hat{C}_3 \quad , \end{aligned}$$

$$\hat{C}_3 = \frac{\sqrt{2k}}{\sqrt{1 + (1 + 4t_\phi^2) \cos^2 \theta_H}} \left\{ \frac{4}{\pi^2 \lambda_n^2} - F_{0,0}^2 \right\}^{-1/2}, \quad (3.33)$$

where $t_\phi \equiv s_\phi/c_\phi$.

Case 3: $\pi^2 \lambda_n^2 z_\pi F_{0,0} F_{1,1} = 2(1 + s_\phi^2) \sin^2 \theta_H$

In this case the mass spectrum depends on θ_H . The lightest mode becomes massless at $\theta_H = 0$ while it acquires a nonzero mass when $\theta_H \neq 0$. The mode is identified with the Z boson. The coefficients in the mode functions are given by

$$\begin{aligned} C_{A,n}^{3L} &= \{c_\phi^2 + \cos \theta_H (1 + s_\phi^2)\} \hat{C}_4, \\ C_{A,n}^{3R} &= \{c_\phi^2 - \cos \theta_H (1 + s_\phi^2)\} \hat{C}_4, \\ C_{A,n}^{\hat{3}} &= -\sqrt{2} (1 + s_\phi^2) \sin \theta_H \frac{F_{1,0}}{F_{1,1}} \hat{C}_4, \\ C_{A,n}^B &= -\frac{2s_\phi c_\phi}{c_\phi^2 + \cos \theta_H (1 + s_\phi^2)} C_{A,n}^{3L}, \\ \hat{C}_4 &= \frac{\sqrt{k}}{\sqrt{2(1 + s_\phi^2) \cos \chi}} \left\{ \frac{4}{\pi^2 \lambda_n^2} + \frac{\pi^2 \lambda_n^2 z_\pi^2 F_{1,0}^2 F_{0,0}^2}{\sin^2 2\chi} - \frac{F_{1,0}^2}{\cos^2 \chi} - \frac{F_{0,0}^2}{\sin^2 \chi} \right\}^{-1/2}, \\ \sin^2 \chi &\equiv \frac{1 + s_\phi^2}{2} \sin^2 \theta_H. \end{aligned} \quad (3.34)$$

Note that the 4×4 matrix in (3.20) reduces to a direct sum of 3×3 and 1×1 matrices and the former is identical to the 3×3 matrix in (3.19) if we set $(s_\phi, c_\phi) = (0, 1)$. Thus the spectrum and the mode functions of the charged sector $(A_\mu^{\pm L}, A_\mu^{\pm R}, A_\mu^{\hat{3}})$ are reproduced from those of $(A_\mu^{3L}, A_\mu^{3R}, A_\mu^{\hat{3}})$ by setting $(s_\phi, c_\phi) = (0, 1)$.

3.2.3. Singlet sector $A_\mu^{\hat{4}}$

Finally there is the singlet sector $A_\mu^{\hat{4}}$. There is no zero mode in this sector. From the normalization condition (3.24), the coefficient is determined as

$$C_{A,n}^{\hat{4}} = \sqrt{2k} \left\{ \frac{4}{\pi^2 \lambda_n^2} - F_{0,1}^2 \right\}^{-1/2}. \quad (3.35)$$

In the gauge sector, there are some classes of K.K. modes whose spectra are independent of θ_H , *i.e.*, the case 1 in the charged sector, the cases 1 and 2 in the neutral sector, and the singlet sector. In all these cases, the mode functions do not have nonzero components for $T^{\hat{a}}$ ($a = 1, 2, 3$, or $\pm, 3$). This means that the modes in these classes do not have nonvanishing couplings to the Higgs field $H = \varphi^{(0)}$, which reflects the θ_H -independence of the mass spectra. (The corresponding coupling constants are expressed like (6.8) in Sect. 6.2.)

3.3. Spectrum and mode functions of gauge scalars

In this subsection the spectrum and mode functions for the extra-dimensional components of gauge potentials, or gauge scalars, are examined. The boundary

conditions for A_z and B_z are given by

$$A_z^{aL} = A_z^{aR} = B_z = 0 \quad , \quad \partial_z \left(\frac{A_z^{\hat{a}}}{z} \right) = 0 \quad (3.36)$$

at both boundaries. The conditions at $z = z_\pi$ determine the ratios between $\alpha_{\varphi,n}$'s and $\beta_{\varphi,n}$'s in Eq.(3.10) so that the mode functions have the following forms.

$$\begin{aligned} \tilde{h}_{\varphi,n}^{aL}(z) &= C_{\varphi,n}^{aL} z F_{0,0}(\lambda_n z, \lambda_n z_\pi) \quad , \\ \tilde{h}_{\varphi,n}^{aR}(z) &= C_{\varphi,n}^{aR} z F_{0,0}(\lambda_n z, \lambda_n z_\pi) \quad , \\ \tilde{h}_{\varphi,n}^{\hat{a}}(z) &= C_{\varphi,n}^{\hat{a}} z F_{0,1}(\lambda_n z, \lambda_n z_\pi) \quad , \\ \tilde{h}_{\varphi,n}^B(z) &= C_{\varphi,n}^B z F_{0,0}(\lambda_n z, \lambda_n z_\pi) \quad . \end{aligned} \quad (3.37)$$

To treat gauge scalars, it is convenient to define

$$A_z^{aL \pm R} \equiv \frac{1}{\sqrt{2}} (A_z^{aL} \pm A_z^{aR}) \quad , \quad (3.38)$$

in terms of which Eq(3.17) can be rewritten as

$$\begin{aligned} \tilde{A}_z^{aL+R} &= A_z^{q^{aL+R}} \quad , \\ \begin{pmatrix} \tilde{A}_z^{aL-R} \\ \tilde{A}_z^{\hat{a}} \end{pmatrix} &= \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} A_z^{q^{aL-R}} \\ A_z^{q^{\hat{a}}} \end{pmatrix} \quad , \\ \tilde{A}_z^{\hat{4}} &= A_z^{q^{\hat{4}}} \quad , \end{aligned} \quad (3.39)$$

where $a = 1, 2, 3$. In contrast to the 4D components A_μ^I and B_μ , the boundary conditions in (3.36) do not mix $\tilde{A}_z^{aL+R}$, $(\tilde{A}_z^{aL-R}, \tilde{A}_z^{\hat{a}})$ and B_z . By making use of (3.37), the boundary conditions at the Planck brane is rewritten as

$$F_{0,0} C_{\varphi,n}^{aL+R} = 0 \quad , \quad (3.40)$$

$$\begin{pmatrix} \cos \theta_H F_{0,0} & -\sin \theta_H F_{0,1} \\ \sin \theta_H F_{1,0} & \cos \theta_H F_{1,1} \end{pmatrix} \begin{pmatrix} C_{\varphi,n}^{aL-R} \\ C_{\varphi,n}^{\hat{a}} \end{pmatrix} = 0 \quad , \quad (3.41)$$

for $a = 1, 2, 3$, and

$$\lambda_n F_{1,1} C_{\varphi,n}^{\hat{4}} = 0 \quad , \quad (3.42)$$

$$F_{0,0} C_{\varphi,n}^B = 0 \quad . \quad (3.43)$$

Here $F_{\alpha,\beta} = F_{\alpha,\beta}(\lambda_n, \lambda_n z_\pi)$. The orthonormal relations are given by

$$\begin{aligned} \int_1^{z_\pi} \frac{kdz}{z} \tilde{h}_{\varphi,n}^{aL+R} \tilde{h}_{\varphi,l}^{aL+R} &= \delta_{n,l} \quad , \\ \int_1^{z_\pi} \frac{kdz}{z} \left\{ \tilde{h}_{\varphi,n}^{aL-R} \tilde{h}_{\varphi,l}^{aL-R} + \tilde{h}_{\varphi,n}^{\hat{a}} \tilde{h}_{\varphi,l}^{\hat{a}} \right\} &= \delta_{n,l} \quad , \end{aligned}$$

$$\int_1^{z_\pi} \frac{k dz}{z} \tilde{h}_{\varphi,n}^{\hat{4}} \tilde{h}_{\varphi,l}^{\hat{4}} = \delta_{n,l} ,$$

$$\int_1^{z_\pi} \frac{k dz}{z} \tilde{h}_{\varphi,n}^B \tilde{h}_{\varphi,l}^B = \delta_{n,l} . \quad (3.44)$$

It follows from the conditions (3.40) and (3.41) that the spectra for the charged sector (3.22) and the neutral sector (3.23) are degenerate. The gauge scalar sector is classified into four cases specified by Eqs.(3.40)-(3.43).

Case 1: Singlet sector I $A_z^{a_{L+R}}$

For the $A_z^{a_{L+R}}$ components the mass spectrum is determined by

$$F_{0,0} = 0 . \quad (3.45)$$

There are no zero-modes and the coefficients $C_{\varphi,n}^{\bar{I}}$ in the mode functions are determined by the normalization condition (3.44).

$$C_{\varphi,n}^{a_{L+R}} = \sqrt{\frac{2}{k}} \left\{ \frac{4}{\pi^2 \lambda_n^2} - F_{1,0}^2 \right\}^{-1/2} ,$$

$$C_{\varphi,n}^{a_{L-R}} = C_{\varphi,n}^{\hat{a}} = C_{\varphi,n}^B = 0 , \quad (3.46)$$

where $a_{L\pm R} = 1, 2, 3$ and $\hat{a} = 1, 2, 3, 4$.

Case 2: Doublet sector $(A_z^{a_{L-R}}, A_z^{\hat{a}})$

For $(A_z^{a_{L-R}}, A_z^{\hat{a}})$ the mass spectrum is determined by

$$F_{0,0} F_{1,1} = \frac{4 \sin^2 \theta_H}{\pi^2 \lambda_n^2 z_\pi} . \quad (3.47)$$

From Eq.(3.41) and the normalization condition (3.44), the mode functions are obtained as

$$C_{\varphi,n}^{a_{L-R}} = \sqrt{\frac{2}{k}} \left\{ \frac{4}{\pi^2 \lambda_n^2} + \frac{\pi^2 \lambda_n^2 z_\pi^2}{\sin^2 2\theta_H} F_{1,0}^2 F_{0,0}^2 - \frac{F_{1,0}^2}{\cos^2 \theta_H} - \frac{F_{0,0}^2}{\sin^2 \theta_H} \right\}^{-1/2} ,$$

$$C_{\varphi,n}^{\hat{a}} = -\tan \theta_H \frac{F_{1,0}}{F_{1,1}} C_{\varphi,n}^{a_{L-R}} ,$$

$$C_{\varphi,n}^{a_{L+R}} = C_{\varphi,n}^B = 0 . \quad (3.48)$$

Case 3: Higgs sector $A_z^{\hat{4}}$

The sector $A_z^{\hat{4}}$ corresponds to the Higgs sector. The spectrum is determined by

$$\lambda_n F_{1,1} = 0 . \quad (3.49)$$

There is a zero-mode, which is identified as the 4D Higgs boson. It acquires a nonvanishing finite mass m_H by quantum effects at the one loop level. It has been

estimated in Refs. 36), 37) that $m_H \sim 0.1\sqrt{\alpha_W} k\pi R m_W / |\sin \theta_H|$, which gives m_H in a range $140 \sim 280$ GeV. The mode functions in this case are given by

$$\begin{aligned}\tilde{h}_{\varphi,0}^{\hat{4}}(z) &= \sqrt{\frac{2}{k(z_\pi^2 - 1)}} z \quad , \\ \tilde{h}_{\varphi,0}^{a_{L\pm R}}(z) &= \tilde{h}_{\varphi,0}^{\hat{a}}(z) = \tilde{h}_{\varphi,0}^B(z) = 0 \quad ,\end{aligned}\tag{3.50}$$

for the zero-mode (the 4D Higgs field), and

$$\begin{aligned}C_{\varphi,n}^{\hat{4}} &= \sqrt{\frac{2}{k}} \left\{ \frac{4}{\pi^2 \lambda_n^2} - F_{0,1}^2 \right\}^{-1/2} \quad , \\ C_{\varphi,n}^{a_{L\pm R}} &= C_{\varphi,n}^{\hat{a}} = C_{\varphi,n}^B = 0 \quad ,\end{aligned}\tag{3.51}$$

for other KK modes ($n \neq 0$).

Case 4: Singlet sector II B_z

The spectrum is the same as in Case 1, but the mode functions are non-vanishing only in the B_z part. They are given by

$$\begin{aligned}C_{\varphi,n}^B &= \sqrt{\frac{2}{k}} \left\{ \frac{4}{\pi^2 \lambda_n^2} - F_{1,0}^2 \right\}^{-1/2} \quad , \\ C_{\varphi,n}^{a_{L\pm R}} &= C_{\varphi,n}^{\hat{a}} = 0,\end{aligned}\tag{3.52}$$

where $a_{L\pm R} = 1, 2, 3$ and $\hat{a} = 1, 2, 3, 4$.

Notice that the spectrum depends on θ_H only in the doublet sector. In other words, the Higgs field couples only to the doublet sector.

§4. Spectrum and mode functions of fermions

Masses of quarks and leptons can originate not only from gauge interactions and bulk kink masses in the fifth dimension, but also from brane interactions with additional fermion fields on the branes. Indeed such additional interactions seem necessary to realize the observed mass spectrum and gauge couplings in the quark and lepton sectors. The main focus in the present paper is gauge-Higgs interactions, and we defer, to a separate paper, detailed discussions about how to construct realistic models. At the moment we merely mention that one can introduce chiral spinor fields χ_R on the Planck brane and χ_L on the TeV brane, which have mixing terms with the bulk fermion Ψ . Let us take $\eta_{0,\pi} = +1$ in (2.6) so that the Z_2 -parities are assigned as Table II for a fermion multiplet in the spinor representation of $SO(5)$. We further suppose that χ_{Ri} ($i = 1, 2$) and χ_L have the same quantum number as Q_{Ri} and q_L , where i denotes the $SU(2)_R$ -doublet index. The Lagrangian in the fermionic sector, then, would be

$$\mathcal{L}_{\text{ferm}} = \sqrt{-G} \left[i\bar{\Psi} \Gamma^N \mathcal{D}_N \Psi - iM_\Psi \varepsilon \bar{\Psi} \Psi \right]$$

Table II. The Z_2 -parities of the fermions. We take the same parity assignment at both boundaries.

q_R	Q_R	q_L	Q_L
even	odd	odd	even

$$\begin{aligned}
& + \sum_{i=1}^2 \{ i \bar{\chi}_{Ri} \gamma^\nu \mathcal{D}_\nu \chi_{Ri} - (i \mu_{Qi} \bar{\chi}_{Ri} Q_{Li} + \text{h.c.}) \} \delta(y) \\
& + \left\{ i \bar{\chi}_L \gamma^\mu \mathcal{D}_\mu \chi_L - (i \mu_q \bar{\chi}_L q_R + \text{h.c.}) \right\} \delta(y - \pi R) \Big] , \quad (4.1)
\end{aligned}$$

where μ_Q and μ_q are brane-mass parameters of mass-dimension 1/2. With these additional parameters a realistic spectrum can be reproduced.

In the subsequent discussions, however, we restrict ourselves to fermions without brane interactions, setting $\mu_Q = \mu_q = 0$ and dropping χ_{Ri} and χ_L . Accordingly the index i of Q_{Li} is suppressed. We describe below how the mass spectrum and gauge couplings are determined, and what kind of potential problems arise in the simplified model.

4.1. General solutions in the bulk

The linearized equations of motion are

$$\begin{aligned}
e^\sigma \gamma^\mu \partial_\mu q_R - (\partial_y - 2\sigma' + M_\Psi \varepsilon) q_L + \frac{ig_A}{2\sqrt{2}} A_y^{c\hat{4}} Q_L &= 0 , \\
e^\sigma \gamma^\mu \partial_\mu Q_R - (\partial_y - 2\sigma' + M_\Psi \varepsilon) Q_L + \frac{ig_A}{2\sqrt{2}} A_y^{c\hat{4}} q_L &= 0 , \\
e^\sigma \gamma^\mu \partial_\mu q_L + (\partial_y - 2\sigma' - M_\Psi \varepsilon) q_R - \frac{ig_A}{2\sqrt{2}} A_y^{c\hat{4}} Q_R &= 0 , \\
e^\sigma \gamma^\mu \partial_\mu Q_L + (\partial_y - 2\sigma' - M_\Psi \varepsilon) Q_R - \frac{ig_A}{2\sqrt{2}} A_y^{c\hat{4}} q_R &= 0 . \quad (4.2)
\end{aligned}$$

From the parity assignment and linearized equations of motion, the boundary conditions for the bulk fermions (q, Q) are determined. In the conformal coordinate $z = e^{\sigma(y)}$, they are written as

$$\begin{aligned}
D_-(c) \hat{q}_R &= 0 , & \hat{q}_L &= 0 , \\
\hat{Q}_R &= 0 , & D_+(c) \hat{Q}_L &= 0 , \quad (4.3)
\end{aligned}$$

at both $z = 1$ and $z = z_\pi$. Here $\hat{q} \equiv z^{-2} q$, $\hat{Q} \equiv z^{-2} Q$, and

$$D_\pm(c) \equiv \pm \frac{d}{dz} + \frac{c}{z} , \quad (4.4)$$

where $c \equiv M_\Psi/k$. We remark that when there are brane interactions with localized fermions, i.e. when $\mu_q, \mu_Q \neq 0$, the above boundary conditions are modified, becoming no longer Dirichlet- nor Neumann-type.

We expand the 5D fermion fields into 4D K.K. modes.

$$\begin{aligned}\hat{q}_R(x, z) &= \sum_n \hat{f}_{R,n}^q(z) \psi_R^{(n)}(x), & \hat{q}_L(x, z) &= \sum_n \hat{f}_{L,n}^q(z) \psi_L^{(n)}(x), \\ \hat{Q}_R(x, z) &= \sum_n \hat{f}_{R,n}^Q(z) \psi_R^{(n)}(x), & \hat{Q}_L(x, z) &= \sum_n \hat{f}_{L,n}^Q(z) \psi_L^{(n)}(x).\end{aligned}\quad (4.5)$$

It follows from (4.2) that equations for an eigenmode with a mass eigenvalue $m_n = k\lambda_n$ are given by

$$\begin{aligned}D_-(c) \hat{f}_{R,n}^q(z) - i \frac{\dot{\theta}}{2} \hat{f}_{R,n}^Q(z) &= \lambda_n \hat{f}_{L,n}^q(z), \\ D_-(c) \hat{f}_{R,n}^Q(z) - i \frac{\dot{\theta}}{2} \hat{f}_{R,n}^q(z) &= \lambda_n \hat{f}_{L,n}^Q(z), \\ D_+(c) \hat{f}_{L,n}^q(z) + i \frac{\dot{\theta}}{2} \hat{f}_{L,n}^Q(z) &= \lambda_n \hat{f}_{R,n}^q(z), \\ D_+(c) \hat{f}_{L,n}^Q(z) + i \frac{\dot{\theta}}{2} \hat{f}_{L,n}^q(z) &= \lambda_n \hat{f}_{R,n}^Q(z).\end{aligned}\quad (4.6)$$

Here $\dot{\theta} \equiv d\theta/dz$ where $\theta(z)$ is defined in (3.16). The orthonormal relations are

$$\begin{aligned}\int_1^{z_\pi} \frac{dz}{k} \left\{ \left(\hat{f}_{R,l}^q \right)^* \hat{f}_{R,n}^q + \left(\hat{f}_{R,l}^Q \right)^* \hat{f}_{R,n}^Q \right\} &= \delta_{n,l} \quad , \\ \int_1^{z_\pi} \frac{dz}{k} \left\{ \left(\hat{f}_{L,l}^q \right)^* \hat{f}_{L,n}^q + \left(\hat{f}_{L,l}^Q \right)^* \hat{f}_{L,n}^Q \right\} &= \delta_{n,l} \quad .\end{aligned}\quad (4.7)$$

In order to solve the mode equations, it is convenient to move to the twisted basis defined in (3.6) and (3.15), in which

$$\begin{pmatrix} \tilde{q} \\ \tilde{Q} \end{pmatrix} \equiv \Omega(z) \begin{pmatrix} \hat{q} \\ \hat{Q} \end{pmatrix} \quad . \quad (4.8)$$

The mode equations are simplified to

$$\begin{aligned}D_-(c) \tilde{f}_{R,n}^q &= \lambda_n \tilde{f}_{L,n}^q, & D_+(c) \tilde{f}_{L,n}^q &= \lambda_n \tilde{f}_{R,n}^q, \\ D_-(c) \tilde{f}_{R,n}^Q &= \lambda_n \tilde{f}_{L,n}^Q, & D_+(c) \tilde{f}_{L,n}^Q &= \lambda_n \tilde{f}_{R,n}^Q,\end{aligned}\quad (4.9)$$

where

$$\begin{pmatrix} \tilde{f}_{R,n}^q \\ \tilde{f}_{R,n}^Q \end{pmatrix} \equiv \begin{pmatrix} c_\theta & is_\theta \\ is_\theta & c_\theta \end{pmatrix} \begin{pmatrix} \hat{f}_{R,n}^q \\ \hat{f}_{R,n}^Q \end{pmatrix}, \quad \begin{pmatrix} \tilde{f}_{L,n}^q \\ \tilde{f}_{L,n}^Q \end{pmatrix} \equiv \begin{pmatrix} c_\theta & is_\theta \\ is_\theta & c_\theta \end{pmatrix} \begin{pmatrix} \hat{f}_{L,n}^q \\ \hat{f}_{L,n}^Q \end{pmatrix}. \quad (4.10)$$

Then (4.5) becomes

$$\begin{aligned}\tilde{q}_R(x, z) &= \sum_n \tilde{f}_{R,n}^q(z) \psi_R^{(n)}(x), & \tilde{q}_L(x, z) &= \sum_n \tilde{f}_{L,n}^q(z) \psi_L^{(n)}(x), \\ \tilde{Q}_R(x, z) &= \sum_n \tilde{f}_{R,n}^Q(z) \psi_R^{(n)}(x), & \tilde{Q}_L(x, z) &= \sum_n \tilde{f}_{L,n}^Q(z) \psi_L^{(n)}(x).\end{aligned}\quad (4.11)$$

The general solutions of Eqs.(4.9) are

$$\begin{aligned}\tilde{f}_{R,n}^q(z) &= z^{\frac{1}{2}} \{a_n^q J_{\alpha-1}(\lambda_n z) + b_n^q Y_{\alpha-1}(\lambda_n z)\}, \\ \tilde{f}_{R,n}^Q(z) &= z^{\frac{1}{2}} \{a_n^Q J_{\alpha-1}(\lambda_n z) + b_n^Q Y_{\alpha-1}(\lambda_n z)\}, \\ \tilde{f}_{L,n}^q(z) &= z^{\frac{1}{2}} \{a_n^q J_{\alpha}(\lambda_n z) + b_n^q Y_{\alpha}(\lambda_n z)\}, \\ \tilde{f}_{L,n}^Q(z) &= z^{\frac{1}{2}} \{a_n^Q J_{\alpha}(\lambda_n z) + b_n^Q Y_{\alpha}(\lambda_n z)\},\end{aligned}\quad (4.12)$$

where $\alpha \equiv (M_{\Psi}/k) + \frac{1}{2}$. The eigenvalues λ_n and the coefficients a_n 's and b_n 's are determined by the boundary conditions (4.3).

4.2. Mass eigenvalues and mode functions

From the conditions (4.3) at $z = z_{\pi}$, the ratios between a_n 's and b_n 's in (4.12) are determined so that the mode functions are written by using the function $F_{\alpha,\beta}(u, v)$ defined in (C.1) as

$$\begin{aligned}\tilde{f}_{R,n}^q(z) &= C_n^q z^{\frac{1}{2}} F_{\alpha-1,\alpha}(\lambda_n z, \lambda_n z_{\pi}), \\ \tilde{f}_{R,n}^Q(z) &= C_n^Q z^{\frac{1}{2}} F_{\alpha-1,\alpha-1}(\lambda_n z, \lambda_n z_{\pi}), \\ \tilde{f}_{L,n}^q(z) &= C_n^q z^{\frac{1}{2}} F_{\alpha,\alpha}(\lambda_n z, \lambda_n z_{\pi}), \\ \tilde{f}_{L,n}^Q(z) &= C_n^Q z^{\frac{1}{2}} F_{\alpha,\alpha-1}(\lambda_n z, \lambda_n z_{\pi}).\end{aligned}\quad (4.13)$$

The eigenvalue λ_n and constants C_n^q and C_n^Q are determined by the remaining boundary conditions in (4.3) at $z = 1$. By making use of (4.13), the two conditions for right-handed components in (4.3) at $z = z_{\pi}$ are rewritten as ^{*)}

$$\begin{pmatrix} c_H F_{\alpha,\alpha} & -i s_H F_{\alpha,\alpha-1} \\ -i s_H F_{\alpha-1,\alpha} & c_H F_{\alpha-1,\alpha-1} \end{pmatrix} \begin{pmatrix} C_n^q \\ C_n^Q \end{pmatrix} = 0. \quad (4.14)$$

Here $F_{\alpha,\beta} = F_{\alpha,\beta}(\lambda_n, \lambda_n z_{\pi})$. For a nontrivial solution to exist, the determinant of the 2×2 matrix in Eq.(4.14) must vanish, which leads to

$$\pi^2 \lambda_n^2 z_{\pi} F_{\alpha-1,\alpha-1} F_{\alpha,\alpha} = 4 s_H^2. \quad (4.15)$$

We have used the last formula in (C.2). This is the equation that determines the mass spectrum $\{\lambda_n\}$. Once λ_n is determined, the corresponding C_n^q and C_n^Q are fixed by (4.14) with the normalization condition (4.7). The result is

$$\begin{aligned}C_n^q &= i t_H \frac{F_{\alpha,\alpha-1}}{F_{\alpha,\alpha}} C_n^Q, \\ C_n^Q &= \sqrt{2k} \left[\frac{4}{\pi^2 \lambda_n^2} + \frac{\pi^2 \lambda_n^2 z_{\pi}^2}{4 s_H^2 c_H^2} F_{\alpha,\alpha-1}^2 F_{\alpha-1,\alpha-1}^2 - \frac{F_{\alpha,\alpha-1}^2}{c_H^2} - \frac{F_{\alpha-1,\alpha-1}^2}{s_H^2} \right]^{-1/2}.\end{aligned}\quad (4.16)$$

^{*)} The remaining conditions in (4.3) at $z = z_{\pi}$ provide the same conditions on (C_n^q, C_n^Q) as (4.14).

§5. Masses and wave functions of light particles

Approximate expressions of the masses and wave functions of light particles such as W , Z , quarks and leptons can be obtained. The mass of the 4D Higgs particle is generated at the quantum level. Its mass is estimated from the effective potential for the Yang-Mills AB phase θ_H .

5.1. Gauge sector

The masses and wave functions of the W and Z bosons have nontrivial θ_H dependence. They belong to the case 2 of the charged sector in Sec. 3.2.1 and the case 3 of the neutral sector in Sec. 3.2.2 whose mass spectra are determined by

$$\begin{aligned}\pi^2 \lambda_n^2 z_\pi F_{0,0} F_{1,1} &= 2 \sin^2 \theta_H, \\ \pi^2 \lambda_n^2 z_\pi F_{0,0} F_{1,1} &= 2(1 + s_\phi^2) \sin^2 \theta_H,\end{aligned}\quad (5.1)$$

respectively. Here $F_{\alpha,\beta} = F_{\alpha,\beta}(\lambda_n, \lambda_n z_\pi)$. These equations are similar to those in the $SU(3)$ model discussed in Ref. 37), but there is an important difference in the numerical factors on the right sides of the two equations above. In the $SU(3)$ model one has $4 \sin^2 \frac{1}{2} \theta_H$ in place of $2 \sin^2 \theta_H$ in the equation determining the W mass.*) Fig. 1 depicts the masses of the KK tower of the W boson as functions of θ_H for $k\pi R = 35, 3.5, 0.35$. Due to the numerical factor mentioned above, the mass spectrum does not approach a linear spectrum in θ_H in the flat limit ($k\pi R \rightarrow 0$) in contrast to the $SU(3)$ model. (See Fig. 1 in Ref. 37).) For $0 < k\pi R \ll 1$,

$$\lambda_n z_\pi \gtrsim \lambda_n \gtrsim \frac{m_{KK}}{k} = \frac{\pi}{z_\pi - 1} \simeq \frac{\pi}{k\pi R} \gg 1 \quad (n \geq 1). \quad (5.2)$$

With (B.3), the left-hand side of (5.1) becomes

$$\pi^2 \lambda_n^2 z_\pi F_{0,0} F_{1,1} \simeq 4 \sin^2(\pi R m_n) \quad (n \geq 1). \quad (5.3)$$

The mass eigenvalues $m_n = k\lambda_n$ become linear functions of θ_H in the flat limit only if the numerical factor in the right-hand side of Eq. (5.1) is 4.**) Therefore the K.K. level-crossing does not occur as θ_H increases from 0 to π in our model even in the flat geometry. This is one of the distinctive properties of the $SO(5) \times U(1)_{B-L}$ model.

As can be seen from Fig. 1, the mass of the lightest mode is much lighter than the K.K. mass scale m_{KK} when the warp factor $z_\pi = e^{k\pi R}$ is large, that is, when $\lambda_0, \lambda_0 z_\pi \ll 1$. Thus one may use (C.4) to obtain an approximate expression for the mass of the lightest mode in each category determined by (5.1) and the corresponding mode function from (3.28) or (3.34). For the W boson the mass is given by

$$m_W = \frac{m_{KK}}{\pi} \sqrt{\frac{1}{k\pi R}} |\sin \theta_H| \left\{ 1 + \mathcal{O}\left(\frac{\pi^2 m_W^2}{m_{KK}^2}\right) \right\} \quad (5.4)$$

*) The masses of the Z boson and its K.K. modes were not discussed in Ref. 37) as the correct value of the Weinberg angle θ_W is not obtained in the $SU(3)$ model.

**) When the numerical factor is 4, it can be easily shown that the mass of the lightest ($n = 0$) mode also becomes a linear function of θ_H in the flat limit.




Fig. 1. The masses of W boson and its K.K. excited states are depicted in the unit of m_{KK} as functions of θ_W for $k\pi R = 35, 3.5$, and 0.35 from the left to the right.

with its mode function

$$\tilde{h}_{A,0}^{\pm L}(z) \simeq \frac{1 + \cos \theta_H}{2\sqrt{\pi R}} \quad , \quad \tilde{h}_{A,0}^{\pm R}(z) \simeq \frac{1 - \cos \theta_H}{2\sqrt{\pi R}} \quad , \quad \tilde{h}_{A,0}^{\pm}(z) \simeq -\frac{\sin \theta_H}{\sqrt{2\pi R}} \left(1 - \frac{z^2}{z_\pi^2}\right) \quad . \quad (5.5)$$

For the Z boson the mass is given by

$$m_Z = \frac{m_{\text{KK}}}{\pi} \sqrt{\frac{1 + s_\phi^2}{k\pi R}} |\sin \theta_H| \left\{ 1 + \mathcal{O}\left(\frac{\pi^2 m_Z^2}{m_{\text{KK}}^2}\right) \right\} \quad (5.6)$$

with its mode function

$$\begin{aligned} \tilde{h}_{A,0}^{3L}(z) &\simeq \frac{c_\phi^2 + \cos \theta_H (1 + s_\phi^2)}{2\sqrt{(1 + s_\phi^2)\pi R}} \quad , \quad \tilde{h}_{A,0}^{3R}(z) \simeq \frac{c_\phi^2 - \cos \theta_H (1 + s_\phi^2)}{2\sqrt{(1 + s_\phi^2)\pi R}} \quad , \\ \tilde{h}_{A,0}^{\hat{3}}(z) &\simeq -\sin \theta_H \sqrt{\frac{1 + s_\phi^2}{2\pi R}} \left(1 - \frac{z^2}{z_\pi^2}\right) \quad , \quad \tilde{h}_{A,0}^B(z) \simeq -\frac{s_\phi c_\phi}{\sqrt{(1 + s_\phi^2)\pi R}} \quad . \end{aligned} \quad (5.7)$$

We stress that m_W and m_Z are not proportional to the VEV of the 4D Higgs field, or θ_H , in contrast to the ordinary Higgs mechanism in four dimensions. The mechanism of mass generation for the 4D gauge bosons in the gauge-Higgs unification scenario involves not only 4D gauge fields and scalar fields in each KK level, but also fields in other KK levels. The lowest mode in each KK tower necessarily mixes with heavy K.K. modes when θ_H acquires a nonzero value. Furthermore, there is mixing with other components of the gauge group at non-vanishing θ_H as well. The spectrum and mixing is such that they become periodic in θ_H with a period 2π .

From (5.4) and (5.6) the Weinberg angle θ_W determined from m_W and m_Z becomes

$$\begin{aligned} \sin^2 \theta_W &\equiv 1 - \frac{m_W^2}{m_Z^2} \\ &\simeq \frac{s_\phi^2}{1 + s_\phi^2} = \frac{g_B^2}{g_A^2 + 2g_B^2} = \frac{g_Y^2}{g_A^2 + g_Y^2} \quad . \end{aligned} \quad (5.8)$$

The approximate equality in the second line is valid to the $\mathcal{O}(0.1\%)$ accuracy for $m_{\text{KK}} = \mathcal{O}(\text{TeV})$. In the last equality the relation $g_Y = g_A g_B / \sqrt{g_A^2 + g_B^2}$ has been

made use of. Note that s_ϕ defined in (2.11) satisfies $s_\phi \simeq \tan \theta_W$. The Weinberg angle θ_W may be determined from the vertices in the neutral current interactions. As we will see in (7.5) below, θ_W determined this way coincides with that in (5.8) to good accuracy. Thus the ρ parameter is nearly one in our model. We remark that the ρ parameter substantially deviates from 1 in the flat limit ($\pi k R \rightarrow 0$) when θ_H is nonvanishing.

According to the classification in the gauge sector, it is convenient to divide the K.K. modes of the charged sector into two classes: $(\tilde{W}_\mu^{(n)}, W_\mu^{(n)})$, corresponding to the cases 1 and 2 in the charged sector. Similarly the K.K. modes of the neutral sector are divided into three classes: $(A_\mu^{\gamma(n)}, \tilde{Z}_\mu^{(n)}, Z_\mu^{(n)})$, corresponding to the cases 1, 2, and 3 in the neutral sector. The W and Z bosons are $W_\mu^{(0)}$ and $Z_\mu^{(0)}$, respectively.

5.2. Fermion sector

The mass spectrum in the fermion sector is determined by Eq.(4.15). For $\lambda_0 z_\pi \ll 1$ we obtain from (4.15) and (C.4)^{37),40)}

$$m_0 = k\lambda_0 \simeq k \left(\frac{\alpha(\alpha-1)}{z_\pi \sinh(\alpha k\pi R) \sinh((\alpha-1)k\pi R)} \right)^{1/2} |s_H|. \quad (5.9)$$

The mode functions for the lightest mode $\psi^{(0)}$ are obtained from Eq.(4.14) with the normalization condition (4.7). For $\alpha > 1$, for example, they are approximately expressed as ^{*})

$$\begin{aligned} \tilde{f}_{R,0}^q(z) &\simeq -i\sqrt{2k\alpha} z_\pi^{-\alpha} z^{\alpha-\frac{1}{2}} \quad , \quad \tilde{f}_{R,0}^Q(z) \simeq s_H c_H \sqrt{2k\alpha} z_\pi^{-\alpha} z^{\frac{3}{2}-\alpha} \quad , \\ \tilde{f}_{L,0}^q(z) &\simeq i s_H \sqrt{2k(\alpha-1)} z^{\frac{1}{2}-\alpha} \quad , \quad \tilde{f}_{L,0}^Q(z) \simeq c_H \sqrt{2k(\alpha-1)} z^{\frac{1}{2}-\alpha} \quad . \end{aligned} \quad (5.10)$$

It follows that the right-handed component $\psi_R^{(0)}$ is localized around the TeV brane while the left-handed component $\psi_L^{(0)}$ around the Planck brane.^{**)} For $\alpha < 0$, on the other hand, $\psi_R^{(0)}$ is localized around the Planck brane while $\psi_L^{(0)}$ around the TeV brane.

With (5.9) the hierarchical fermion mass spectrum can be reproduced by choosing α in an $\mathcal{O}(1)$ range.³⁷⁾ However there is a serious problem in this scenario as pointed out in Ref. 1). The gauge couplings to the W and Z bosons deviate from the observed values for at least one chiral component of the fermion when θ_H is nonzero. This is due to the fact that the wave function of one of the chiral components is inevitably localized around the TeV brane where the mode functions for the W and Z bosons deviate from the constant values (see (5.5) and (5.7)).

This problem can be avoided by introducing boundary fields and turning on boundary mass terms with bulk fermions as described with the action (4.1). There

^{*}) The subleading terms in $\tilde{f}_{L,0}^q(z)$ and $\tilde{f}_{R,0}^Q(z)$ dropped in (5.10) become comparable to the leading terms at $z = z_\pi$, but they remain suppressed in the bulk.

^{**)} $\tilde{f}_{R,0}^Q(z)$ can be localized at the Planck brane if $\alpha > 3/2$, but it is exponentially suppressed compared to $\tilde{f}_{R,0}^q(z)$ for $z > 1$ in such a case.

are two origins for fermion masses; the Yang-Mills AB phase θ_H and boundary masses μ_q, μ_Q . It can be shown that for $\alpha > 1$ and $z_\pi^{2(1-\alpha)} \ll (\mu_Q^2/k)z_\pi^2 \ll \mu_q^2/k \simeq \mathcal{O}(1)$ the lightest mass becomes $m_0 \simeq \sqrt{k(\alpha-1)}\mu_Q$ and its wave functions are suppressed at the TeV brane for both L and R chiral components. Further $SU(2)_R$ is broken at the Planck brane so that μ_Q can be chosen to be different for the upper and the lower components of Q_L . Thus one can realize the observed fermion masses with appropriate μ_Q . The detailed discussions will be given in a separate paper.

§6. Gauge-Higgs self-couplings

Once wave functions of the W and Z bosons and the Higgs boson are determined, the effective four-dimensional couplings among them can be calculated as overlap integrals in the fifth dimension. As θ_H becomes nonvanishing, the form of wave functions substantially changes with large mixing so that the effective four-dimensional couplings are expected to have nontrivial θ_H -dependence in general. This behavior of the couplings provides crucial tests for the gauge-Higgs unification scenario.

The WWZ coupling has been indirectly measured in the LEP2 experiment of W pair production. The standard model fits the data well within a few percents so that deviation from the value in the standard mode could rule out the model under consideration. As we shall see below, the WWZ coupling in the gauge-Higgs unification in the Randall-Sundrum warped spacetime remains almost universal as in the standard model. In the gauge-Higgs unification in flat spacetime, however, substantial deviation results.

Important predictions from the gauge-Higgs unification scenario are obtained for the WWH , ZZH , $WWHH$ and $ZZHH$ couplings where H stands for the Higgs boson. These couplings are suppressed compared with those in the standard model. This gives a crucial test to be performed at LHC in the coming years.

6.1. WWZ coupling

The self-couplings among W , Z and Higgs bosons are determined from the interaction terms in the twisted basis

$$\int_{z_0}^{z_\pi} \frac{dz}{kz} \left\{ ig_A \eta^{\mu\rho} \eta^{\nu\sigma} \text{tr}(\partial_\mu \tilde{A}_\nu - \partial_\nu \tilde{A}_\mu) [\tilde{A}_\rho, \tilde{A}_\sigma] + 2ig_A k^2 \eta^{\mu\nu} \text{tr}(\partial_\mu \tilde{A}_z - \partial_z \tilde{A}_\mu) [\tilde{A}_\nu, \tilde{A}_z] \right. \\ \left. + \frac{1}{2} g_A^2 \eta^{\mu\rho} \eta^{\nu\sigma} \text{tr}[\tilde{A}_\mu, \tilde{A}_\nu] [\tilde{A}_\rho, \tilde{A}_\sigma] + g_A^2 k^2 \eta^{\mu\nu} \text{tr}[\tilde{A}_\mu, \tilde{A}_z] [\tilde{A}_\nu, \tilde{A}_z] \right\} \quad (6.1)$$

by inserting the wave functions of W , Z and H . The relevant part of the expansion of the gauge fields is

$$\tilde{A}_\mu = W_\mu^{(0)}(x) \left\{ \tilde{h}_W^{\pm L}(z) T^{-L} + \tilde{h}_W^{\pm R}(z) T^{-R} + \tilde{h}_W^{\pm}(z) T^{\hat{}} \right\} \\ + W_\mu^{(0)\dagger}(x) \left\{ \tilde{h}_W^{\pm L}(z) T^{+L} + \tilde{h}_W^{\pm R}(z) T^{+R} + \tilde{h}_W^{\pm}(z) T^{\hat{}} \right\} \\ + Z_\mu^{(0)}(x) \left\{ \tilde{h}_Z^{3L}(z) T^{3L} + \tilde{h}_Z^{3R}(z) T^{3R} + \tilde{h}_Z^{\hat{3}}(z) T^{\hat{3}} \right\} ,$$

$$\tilde{A}_z = H^{(0)}(x)\tilde{h}_H^4(z)T^4 . \quad (6.2)$$

Here $T^{\pm L} = (T^{1L} \pm iT^{2L})/\sqrt{2}$ etc.. The W wave functions $\tilde{h}_W^{\pm R,L}(z) = \tilde{h}_{A,0}^{\pm R,L}(z)$ and $\tilde{h}_W^{\pm}(z) = \tilde{h}_{A,0}^{\pm}(z)$ are given by (3.18) with (3.28). The Z wave functions $\tilde{h}_Z^{3R,L}(z) = \tilde{h}_{A,0}^{3R,L}(z)$ and $\tilde{h}_Z^3(z) = \tilde{h}_{A,0}^3(z)$ are given by (3.18) with (3.34). The Higgs wave function $\tilde{h}_H^4(z) = \tilde{h}_{\varphi,0}^4(z)$ is given by (3.50).

The WWZ coupling is evaluated by inserting (6.2) into the first term in (6.1) and integrating over z . The result is

$$\begin{aligned} \mathcal{L}_{WWZ}^{(4)} = ig_{WWZ} \Big\{ & (\partial_\mu W_\nu^{(0)\dagger} - \partial_\nu W_\mu^{(0)\dagger}) W^{(0)\mu} Z^{(0)\nu} - (\partial_\mu W_\nu^{(0)} - \partial_\nu W_\mu^{(0)}) W^{(0)\dagger\mu} Z^{(0)\nu} \\ & + W_\mu^{(0)\dagger} W_\nu^{(0)} \left(\partial^\mu Z^{(0)\nu} - \partial^\nu Z^{(0)\mu} \right) \Big\} \end{aligned} \quad (6.3)$$

where the coupling g_{WWZ} is expressed as overlap integrals

$$\begin{aligned} g_{WWZ} = g_A \int_1^{z_\pi} \frac{dz}{kz} \Big[& \tilde{h}_Z^{3L} \left\{ \left(\tilde{h}_W^{\pm L} \right)^2 + \frac{1}{2} \left(\tilde{h}_W^{\pm} \right)^2 \right\} + \tilde{h}_Z^{3R} \left\{ \left(\tilde{h}_W^{\pm R} \right)^2 + \frac{1}{2} \left(\tilde{h}_W^{\pm} \right)^2 \right\} \\ & + \tilde{h}_Z^3 \tilde{h}_W^{\pm} \left(\tilde{h}_W^{\pm L} + \tilde{h}_W^{\pm R} \right) \Big] . \end{aligned} \quad (6.4)$$

Note that given θ_H and $k\pi R$, m_W determines k . $k = O(M_{pl})$ corresponds to $k\pi R \sim 35$. With these parameters we have exact wave functions as summarized in (3.18), (3.28) and (3.34). When $k\pi R \sim 35$ and the warp factor is large $e^{k\pi R} \gg 1$, the approximate expressions for the wave functions of W and Z given by (5.5) and (5.7) can be employed to find

$$g_{WWZ} \simeq \frac{g_A}{\sqrt{(1 + s_\phi^2)\pi R}} \simeq g \cos \theta_W . \quad (6.5)$$

Here a dimensionless coupling g is defined as

$$g \equiv \frac{g_A}{\sqrt{\pi R}} , \quad (6.6)$$

which is the 4D $SU(2)_L$ gauge coupling at $\theta_H = 0$. In the last equality in (6.5), (5.8) has been made use of. We have neglected corrections suppressed by a factor of $(k\pi R)^{-1} \simeq 1/35$ in conformity with the approximation employed in deriving Eqs.(5.4)-(5.7).

These couplings (6.5) have the same values as those in the standard model. Although the wave functions vary substantially as θ_H , the triple gauge coupling WWZ remains almost universal. The statement can be strengthened by numerically integrating (6.4) with exact wave functions. In Table III the values of the ratio of the trilinear couplings (6.4) to $g_A/\sqrt{(1 + s_\phi^2)\pi R}$ are shown for various values of θ_H and $k\pi R$. It is clearly seen that the relation (6.5) hold to extreme accuracy when the warp factor $e^{k\pi R}$ is large. This is very important in the light of the experimental

Table III. The values of a ratio of $g_{WWZ}^{(1)} (= g_{WWZ}^{(2)})$ to $g_A/\sqrt{(1+s_\phi^2)\pi R}$ for $\theta_H = \pi/10, \pi/4, \pi/2$ and $k\pi R = 35, 3.5, 0.35$.

$k\pi R \backslash \theta_H$	$\pi/10$	$\pi/4$	$\pi/2$
35	0.9999987	0.999964	0.99985
3.5	0.9999078	0.996993	0.98460
0.35	0.9994990	0.979458	0.83378

fact that W pair production rate measured at LEP2 is consistent with the WWZ coupling in the standard model. In the flat spacetime limit ($k\pi R \rightarrow 0$), however, substantial deviation from the standard model results with moderate θ_H . When $\theta_H = O(1)$ gauge-Higgs unification in flat spacetime contradicts with the data for the WWZ coupling.

As can be seen in the overlap integrals in (6.4), weight of wave functions is best measured in the $y = k^{-1} \ln z$ coordinate. The wave functions of the gauge bosons in the warped spacetime in the y coordinate remain almost constants except in a tiny region near $y = \pi R$ even at $\theta_H \neq 0$. (See (5.5) and (5.7).) In the flat spacetime, however, wave functions are deformed substantially at $\theta_H \neq 0$ to have nontrivial y -dependence in the entire region, which contributes to substantial deviation in the WWZ coupling from the standard model.

6.2. WWH and ZZH couplings

A striking prediction of the gauge-Higgs unification scenario is obtained for the Higgs couplings to W and Z . Let us consider the WWH and ZZH couplings. The 4D Higgs field $H = \varphi^{(0)}$ is a part of the 5D gauge fields so that the Higgs couplings to W and Z are completely determined by the gauge principle once θ_H is given. Unlike the WWZ coupling, substantial deviation from the standard model is predicted.

Indeed, by inserting (6.2) into the second term in (6.1), one finds that

$$\mathcal{L}^{(4)} = \lambda_{WWH} H^{(0)} W^{(0)\mu\dagger} W_\mu^{(0)} + \frac{1}{2} \lambda_{ZZH} H^{(0)} Z^{(0)\mu} Z_\mu^{(0)} + \dots, \quad (6.7)$$

where

$$\begin{aligned} \lambda_{WWH} &= g_A k \int_1^{z_\pi} \frac{dz}{z} \tilde{h}_H^4 \left\{ \tilde{h}_W^\pm \partial_z \left(\tilde{h}_W^{\pm L} - \tilde{h}_W^{\pm R} \right) - \partial_z \tilde{h}_W^\pm \left(\tilde{h}_W^{\pm L} - \tilde{h}_W^{\pm R} \right) \right\}, \\ \lambda_{ZZH} &= g_A k \int_1^{z_\pi} \frac{dz}{z} \tilde{h}_H^4 \left\{ \tilde{h}_Z^3 \partial_z \left(\tilde{h}_Z^{3L} - \tilde{h}_Z^{3R} \right) - \partial_z \tilde{h}_Z^3 \left(\tilde{h}_Z^{3L} - \tilde{h}_Z^{3R} \right) \right\}. \end{aligned} \quad (6.8)$$

Recall that the wave function of the Higgs field, $\tilde{h}_H^4(z) \propto z = e^{ky}$, is localized near the TeV brane at $z = z_\pi$ when evaluated in the y -coordinate relevant in the integrals (6.8). The behavior of the wave functions of W and Z bosons near the TeV brane sensitively depends on θ_H so that nontrivial θ_H dependence is expected for the WWH and ZZH couplings.

The integrals in (6.8) can be evaluated in a closed form. Consider λ_{WWH} . Inserting the wave functions into (6.8) and making use of the identity $\partial_z \{ z F_{1,\beta}(\lambda z, \lambda z_\pi) \} =$

$\lambda z F_{0,\beta}(\lambda z, \lambda z_\pi)$ and the last relation in (C·2), one finds

$$\begin{aligned}\lambda_{WWH} &= \frac{4g_A k^2 C_W^\pm (C_W^{\pm L} - C_W^{\pm R})}{\pi^2 m_W z_\pi} \int_1^{z_\pi} dz \tilde{h}_H^4(z) \\ &= \frac{2g_A k \sqrt{2k(z_\pi^2 - 1)}}{\pi^2 m_W z_\pi} C_W^\pm (C_W^{\pm L} - C_W^{\pm R}) .\end{aligned}\quad (6.9)$$

The coefficients $C_W^{\pm L,R} = C_{A,0}^{\pm L,R}$ and $C_W^\pm = C_{A,0}^\pm$ are given by (3·28). Similarly for λ_{ZZH} one finds

$$\lambda_{ZZH} = \frac{2g_A k \sqrt{2k(z_\pi^2 - 1)}}{\pi^2 m_Z z_\pi} C_Z^\pm (C_Z^{\pm L} - C_Z^{\pm R}) , \quad (6.10)$$

where the coefficients $C_Z^{\pm L,R} = C_{A,0}^{\pm L,R}$ and $C_Z^\pm = C_{A,0}^\pm$ are given by (3·34).

The formulas (6·9) and (6·10) are exact. They are fairly well evaluated with the approximate formulas (5·4)-(5·7), leading to $\hat{C}_2 \sim \sqrt{\pi} m_W / 4k\sqrt{R}$ for W and $\hat{C}_4 \sim \sqrt{\pi} m_Z / 4k\sqrt{(1 + s_\phi^2)R}$ for Z . Insertion of these gives

$$\begin{aligned}\lambda_{WWH} &\simeq \frac{g_A \sqrt{k}}{\pi R z_\pi} \sin \theta_H \cos \theta_H \simeq g m_W \cdot p_H |\cos \theta_H| , \\ \lambda_{ZZH} &\simeq \frac{g_A \sqrt{k}(1 + s_\phi^2)}{\pi R z_\pi} \sin \theta_H \cos \theta_H \simeq \frac{g m_Z}{\cos \theta_W} \cdot p_H |\cos \theta_H| ,\end{aligned}\quad (6.11)$$

where g is defined in (6·6) and $p_H \equiv \text{sgn}(\tan \theta_H)$. Both λ_{WWH} and λ_{ZZH} are suppressed by a factor $\cos \theta_H$ compared with the corresponding couplings in the standard model. Unless θ_H is very small, this gives substantial suppression which can be checked in the coming experiments at LHC. This is a generic prediction in the gauge-Higgs unification in the warped spacetime. It does not depend on the details of the model such as fermion content and couplings.

6.3. Vanishing $WWH^{(n)}$ and $ZZH^{(n)}$ couplings

There is a Kaluza-Klein (KK) tower of the 4D Higgs field. The 4D Higgs field is a part of the fifth dimensional component of gauge potentials. It is associated with the Yang-Mills AB phase along the fifth dimension, and is a physical degree of freedom. Its KK excited states, however, are unphysical. In the unitary gauge they are eliminated to be absorbed by the KK excited states of the four-dimensional gauge fields. One may wonder if these KK excited states of the 4D Higgs field, $H^{(n)}$, have nontrivial couplings to W and Z .

The $WWH^{(n)}$ and $ZZH^{(n)}$ couplings are evaluated in the same manner as the WWH and ZZH couplings in Sec. 6.2. They are given by

$$\mathcal{L}^{(4)} = \lambda_{WWH^{(n)}} H^{(n)} W^{(0)\mu \dagger} W_\mu^{(0)} + \frac{1}{2} \lambda_{ZZH^{(n)}} H^{(n)} Z^{(0)\mu} Z_\mu^{(0)} + \cdots , \quad (6.12)$$

where

$$\lambda_{WWH^{(n)}} = g_A k \int_1^{z_\pi} \frac{dz}{z} \tilde{h}_{H^{(n)}}^4 \left\{ \tilde{h}_W^\pm \partial_z \left(\tilde{h}_W^{\pm L} - \tilde{h}_W^{\pm R} \right) - \partial_z \tilde{h}_W^\pm \left(\tilde{h}_W^{\pm L} - \tilde{h}_W^{\pm R} \right) \right\} ,$$

$$\lambda_{ZZH^{(n)}} = g_A k \int_1^{z_\pi} \frac{dz}{z} \tilde{h}_{H^{(n)}}^4 \left\{ \tilde{h}_Z^3 \partial_z (\tilde{h}_Z^{3L} - \tilde{h}_Z^{3R}) - \partial_z \tilde{h}_Z^3 (\tilde{h}_Z^{3L} - \tilde{h}_Z^{3R}) \right\} . \quad (6.13)$$

We first recall that the spectrum $m_n^H = k\lambda_n^H$ for $H^{(n)}$ ($n \geq 1$) is determined by the equation $F_{1,1}(\lambda_n^H, \lambda_n^H z_\pi) = 0$. The wave function is given by $\tilde{h}_{H^{(n)}}^4(z) = C_{\varphi,n}^4 z F_{0,1}(\lambda_n^H z, \lambda_n^H z_\pi)$ where $C_{\varphi,n}^4$ is given by (3.51). $\lambda_{WWH^{(n)}}$ is evaluated as λ_{WWH} in the previous subsection. A similar expression to the first line in Eq. (6.9) is obtained where $\tilde{h}_H^4$ is replaced by $\tilde{h}_{H^{(n)}}^4$. Inserting the wave function, one finds

$$\begin{aligned} \lambda_{WWH^{(n)}} &= \frac{4g_A k^2 C_{\varphi,n}^4 C_W^\pm (C_W^{\pm L} - C_W^{\pm R})}{\pi^2 m_W z_\pi} \int_1^{z_\pi} dz z F_{0,1}(\lambda_n^H z, \lambda_n^H z_\pi) \\ &= \frac{4g_A k^2 C_{\varphi,n}^4 C_W^\pm (C_W^{\pm L} - C_W^{\pm R})}{\pi^2 m_W z_\pi} \frac{-1}{\lambda_n^H} F_{1,1}(\lambda_n^H, \lambda_n^H z_\pi) \\ &= 0 . \end{aligned} \quad (6.14)$$

Similarly one finds that $\lambda_{ZZH^{(n)}} = 0$. This proves that the $WWH^{(n)}$ and $ZZH^{(n)}$ couplings identically vanish.

6.4. W^4 and $W^2 Z^2$ couplings

The four-dimensional gauge couplings $WWW^\dagger W^\dagger$ and $WW^\dagger ZZ$ are evaluated from the third term in (6.1). One finds that

$$\begin{aligned} \mathcal{L}^{(4)} &= \frac{1}{2} g_{WWW}^2 \left\{ W_\mu^{(0)\dagger} W^{(0)\dagger\mu} W_\nu^{(0)} W^{(0)\nu} - (W_\mu^{(0)\dagger} W^{(0)\mu})^2 \right\} \\ &\quad + g_{WWZZ}^2 \left\{ W_\mu^{(0)} Z^{(0)\mu} W_\nu^{(0)\dagger} Z^{(0)\nu} - W_\mu^{(0)} W^{(0)\dagger\mu} Z_\nu^{(0)} Z^{(0)\nu} \right\} \end{aligned} \quad (6.15)$$

where the couplings g_{WWW}^2 and g_{WWZZ}^2 are expressed as overlap integrals

$$\begin{aligned} g_{WWW}^2 &= g_A^2 \int_1^{z_\pi} \frac{dz}{kz} \left[\left\{ (\tilde{h}_W^{\pm L})^2 + \frac{1}{2} (\tilde{h}_W^{\pm})^2 \right\}^2 + \left\{ (\tilde{h}_W^{\pm R})^2 + \frac{1}{2} (\tilde{h}_W^{\pm})^2 \right\}^2 \right. \\ &\quad \left. + (\tilde{h}_W^{\pm})^2 (\tilde{h}_W^{\pm L} + \tilde{h}_W^{\pm R})^2 \right] , \\ g_{WWZZ}^2 &= g_A^2 \int_1^{z_\pi} \frac{dz}{kz} \left[\left\{ \tilde{h}_W^{\pm L} \tilde{h}_Z^{3L} + \frac{1}{2} \tilde{h}_W^{\pm} \tilde{h}_Z^3 \right\}^2 + \left\{ \tilde{h}_W^{\pm R} \tilde{h}_Z^{3R} + \frac{1}{2} \tilde{h}_W^{\pm} \tilde{h}_Z^3 \right\}^2 \right. \\ &\quad \left. + \frac{1}{4} \left\{ (\tilde{h}_W^{\pm L} + \tilde{h}_W^{\pm R}) \tilde{h}_Z^3 + \tilde{h}_W^{\pm} (\tilde{h}_Z^{3L} + \tilde{h}_Z^{3R}) \right\}^2 \right] . \end{aligned} \quad (6.16)$$

Inserting (5.5) and (5.7) into (6.16), one finds

$$g_{WWW}^2 \simeq \frac{g_A^2}{\pi R} = g^2 ,$$

$$g_{WWZZ}^2 \simeq \frac{g_A^2}{\pi R(1+s_\phi^2)} = g^2 \cos^2 \theta_W \quad , \quad (6.17)$$

which coincide with couplings in the standard model.

The quartic gauge couplings are approximately independent of θ_H in the gauge-Higgs unification in the warped spacetime, while in flat spacetime they deviate from the values in the standard model. The approximate universality of the gauge couplings in the warped spacetime is guaranteed by the approximately uniform distribution of the wave functions of the gauge bosons in the entire fifth dimension except for the tiny vicinity of the TeV brane.

6.5. $WWHH$ and $ZZHH$ couplings

The four-dimensional gauge-Higgs couplings $WW^\dagger HH$ and $ZZHH$ are evaluated from the fourth term in (6.1). One finds that

$$\mathcal{L}^{(4)} = -\frac{1}{4} \lambda_{WWHH}^2 W_\mu^{(0)\dagger} W^{(0)\mu} H^{(0)} H^{(0)} - \frac{1}{8} \lambda_{ZZHH}^2 Z_\mu^{(0)} Z^{(0)\mu} H^{(0)} H^{(0)} \quad (6.18)$$

where the couplings λ_{WWHH}^2 and λ_{ZZHH}^2 are expressed as overlap integrals

$$\begin{aligned} \lambda_{WWHH}^2 &= k g_A^2 \int_1^{z_\pi} \frac{dz}{z} (\tilde{h}_H^4)^2 \left\{ (\tilde{h}_W^{\pm L} - \tilde{h}_W^{\pm R})^2 + 2(\tilde{h}_W^\pm)^2 \right\} \quad , \\ \lambda_{ZZHH}^2 &= k g_A^2 \int_1^{z_\pi} \frac{dz}{z} (\tilde{h}_H^4)^2 \left\{ (\tilde{h}_Z^{3L} - \tilde{h}_Z^{3R})^2 + 2(\tilde{h}_Z^3)^2 \right\} \quad . \end{aligned} \quad (6.19)$$

As in the case of the cubic couplings λ_{WZH} and λ_{ZZH} , the Higgs wave function $\tilde{h}_{\varphi,0}^4$ is localized near the TeV brane so that the overlap integrals suffer from nontrivial θ_H dependence. With (5.5) and (5.7) inserted, the integrals in (6.19) are evaluated to be

$$\begin{aligned} \lambda_{WWHH}^2 &\simeq \frac{g_A^2}{\pi R} \left(1 - \frac{2}{3} \sin^2 \theta_H \right) = g^2 \left(1 - \frac{2}{3} \sin^2 \theta_H \right) \quad , \\ \lambda_{ZZHH}^2 &\simeq \frac{g_A^2(1+s_\phi^2)}{\pi R} \left(1 - \frac{2}{3} \sin^2 \theta_H \right) \simeq \frac{g^2}{\cos^2 \theta_W} \left(1 - \frac{2}{3} \sin^2 \theta_H \right) \quad . \end{aligned} \quad (6.20)$$

Compared with the values in the standard model, these couplings are suppressed by a factor $(1 - \frac{2}{3} \sin^2 \theta_H)$.

One comment is in order. The couplings defined in (6.19) and (6.20) are to be called as the bare $WW^\dagger HH$ and $ZZHH$ couplings. In the effective theory at low energies where all heavy modes are integrated out, the effective $WW^\dagger HH$ and $ZZHH$ couplings contain contributions coming from tree diagrams involving $W^{(n)}$ and $Z^{(n)}$ as intermediate states. Their contributions may not be negligible, and need careful examination.

The suppression of the bare Higgs couplings is a generic feature of the gauge-Higgs unification, and should be used for testing the scenario by experiments.

§7. Gauge couplings of fermions

Inserting (3·9) and (4·11) into

$$S_{\text{gc}} = \int d^4x \int_1^{z_\pi} \frac{dz}{k} \left\{ g_A \bar{\tilde{\Psi}} \gamma^\mu \tilde{A}_\mu \tilde{\Psi} + \frac{g_B}{2} \bar{\tilde{\Psi}} \gamma^\mu B_\mu \mathcal{Q}_{\text{B-L}} \tilde{\Psi} \right\} , \quad (7.1)$$

we obtain

$$\begin{aligned} \mathcal{L}_{\text{gc}}^{(4)} = & \sum_n W_\mu^{(n)} \left\{ \frac{g_L^{W(n)}}{\sqrt{2}} \bar{\psi}_{L2}^{(0)} \gamma^\mu \psi_{L1}^{(0)} + \frac{g_R^{W(n)}}{\sqrt{2}} \bar{\psi}_{R2}^{(0)} \gamma^\mu \psi_{R1}^{(0)} + \text{h.c.} \right\} \\ & + \sum_n Z_\mu^{(n)} \sum_{i=1}^2 \left\{ g_{Li}^{Z(n)} \bar{\psi}_{Li}^{(0)} \gamma^\mu \psi_{Li}^{(0)} + g_{Ri}^{Z(n)} \bar{\psi}_{Ri}^{(0)} \gamma^\mu \psi_{Ri}^{(0)} \right\} \\ & + \sum_n A_\mu^{\gamma(n)} \sum_{i=1}^2 \left\{ g_{Li}^{\gamma(n)} \bar{\psi}_{Li}^{(0)} \gamma^\mu \psi_{Li}^{(0)} + g_{Ri}^{\gamma(n)} \bar{\psi}_{Ri}^{(0)} \gamma^\mu \psi_{Ri}^{(0)} \right\} + \dots , \end{aligned} \quad (7.2)$$

where the ellipsis denotes terms involving the massive K.K. modes of the fermions. The 4D gauge couplings are given by

$$\begin{aligned} g_L^{W(n)} \equiv & g_A \int_1^{z_\pi} \frac{dz}{k} \left\{ |\tilde{f}_{L,0}^q|^2 \tilde{h}_{W(n)}^{\pm L} + |\tilde{f}_{L,0}^Q|^2 \tilde{h}_{W(n)}^{\pm R} + \sqrt{2} \text{Im} \left\{ \left(\tilde{f}_{L,0}^Q \right)^* \tilde{h}_{W(n)}^{\pm} \tilde{f}_{L,0}^q \right\} \right\} , \\ g_{Li}^{\gamma, Z(n)} \equiv & (-)^{i-1} \frac{g_A}{2} \int_1^{z_\pi} \frac{dz}{k} \left[|\tilde{f}_{L,0}^q|^2 \tilde{h}_{\gamma, Z(n)}^{3L} + |\tilde{f}_{L,0}^Q|^2 \tilde{h}_{\gamma, Z(n)}^{3R} \right. \\ & \left. + \sqrt{2} \text{Im} \left\{ \left(\tilde{f}_{L,0}^Q \right)^* \tilde{h}_{\gamma, Z(n)}^3 \tilde{f}_{L,0}^q \right\} \right] \\ & + \frac{g_B q_{\text{B-L}}}{2} \int_1^{z_\pi} \frac{dz}{k} \tilde{h}_{\gamma, Z(n)}^B \left\{ |\tilde{f}_{L,0}^q|^2 + |\tilde{f}_{L,0}^Q|^2 \right\} , \end{aligned} \quad (7.3)$$

where $q_{\text{B-L}}$ is an eigenvalue of $\mathcal{Q}_{\text{B-L}}$. The index $i = 1, 2$ denotes the $SU(2)_R$ -doublet index. The same expressions hold for the right-handed (R) components where L is replaced by R . The wave functions $\tilde{h}_{\gamma(n)}^{3L}$, $\tilde{h}_{Z(n)}^{3L}$ etc. are given by (3·30), (3·31) and by (3·34), respectively. In a simplified model without boundary mass terms there results nontrivial couplings of right-handed fermions to W bosons.

From the approximate expressions of the mode functions (5·5), (5·7), and (5·10), the 4D gauge couplings are found to be

$$g_L^{W(0)} \simeq \frac{g_A}{\sqrt{\pi R}} = g , \quad (7.4)$$

$$g_L^{Z(0)} \simeq \frac{(-1)^{i-1} g_A - g_B q_{\text{B-L}} s_\phi c_\phi}{2\sqrt{(1 + s_\phi^2)\pi R}} \simeq \frac{g}{\cos \theta_W} \left\{ \frac{(-1)^{i-1}}{2} - q_{\text{EM}} \sin^2 \theta_W \right\} , \quad (7.5)$$

$$g_i^{\gamma(0)} = e q_{\text{EM}} , \quad (7.6)$$

where $e \equiv g_A \sin \theta_W / \sqrt{\pi R} = g \sin \theta_W$ is the $U(1)_{EM}$ gauge coupling constant and $q_{EM} \equiv \{(-1)^{i-1} + q_{B-L}\} / 2$ is the electromagnetic charge. The relation (5.8) has been made use of in the second equality in (7.5). Note that Eqs.(7.4), (7.5) and (7.6) agree with the counterparts in the standard model. Rigorously speaking, the couplings $g_L^{W(0)}$ and $g_L^{Z(0)}$ have small dependence on the parameter α , which leads to tiny violation of the universality in weak interactions as discussed in Ref. 37). It was found that there results violation of the μ - e universality of $O(10^{-8})$, which is well in the experimental bound.

$g_R^{W(0)}$ and $g_R^{Z(0)}$ for a multiplet Ψ , however, substantially deviate from the standard model values. For instance, one finds $g_R^{W(0)} \simeq g(1 - \cos \theta_H)/2$. This is because the mode functions of the right-handed fermions are localized near the TeV brane for $\alpha > 1$. Since K.K. excited states are also localized near the TeV brane, the mixing with K.K. excited states becomes strong, leading to the deviation. The problem can be avoided by introducing boundary fields $\chi_{R,L}$ with boundary mass mixing with bulk fermions, *i.e.*, $\mu_q, \mu_Q \neq 0$ as in the action (4.1). It can be arranged such that right-handed components mainly consist of boundary fields on the Planck brane so that the deviation of the gauge couplings from the standard model become small enough.

§8. Summary

Gauge-Higgs unification in warped spacetime has many attractive features. It identifies the 4D Higgs field with a zero mode of the extra-dimensional components of the gauge fields, or the Yang-Mills AB phase in the extra dimensions. The Higgs couplings are determined by the gauge principle and the structure of background spacetime.

Although the wave functions of the W and Z bosons substantially vary as θ_H , the WWZ , $WWWW$, and $WWZZ$ couplings in the warped spacetime remain nearly the same as in the standard model. However, these couplings considerably deviate from the standard model in the flat spacetime, thus contradicting the LEP2 data on the W pair production. These stem from the fact that the wave functions of the gauge bosons in the warped spacetime remain almost constants except in a tiny region near $y = \pi R$ even at $\theta_H \neq 0$ while they are deformed substantially to have nontrivial y -dependence in the entire region in the flat spacetime. The warped spacetime saves the universality of the gauge couplings.

The important deviation from the standard model shows up in the Higgs couplings. We have shown that the WWH and ZZH couplings are suppressed by a factor $\cos \theta_H$ compared with those in the standard model whereas the bare $WWHH$ and $ZZHH$ couplings are suppressed by a factor $1 - \frac{2}{3} \sin^2 \theta_H$. The precise content of matter fields affects the location of the global minimum of the effective potential $V_{\text{eff}}(\theta_H)$. Once the value of θ_H is determined, the wave functions of W , Z , and H are fixed as functions of θ_H . Hence the suppression of the Higgs couplings to W and Z is a generic feature of the gauge-Higgs unification, and is independent of the details of the model. It can be used to test the scenario. A similar suppression of these Higgs

couplings are recently discussed in Ref. 43) with detailed phenomenological studies in the context of models where the Higgs emerges from a strongly-interacting sector as a pseudo-Goldstone boson, which are closely related to the gauge-Higgs unification models. The phenomenological study of our results such as detailed detectability at LHC/ILC is an important issue and need to be investigated.

As briefly discussed in the present paper, additional brane interactions are necessary to have a realistic spectrum and gauge couplings of fermions. Wave functions of fermions sensitively depend on such interactions, and so do Yukawa couplings. In Ref. 37) it is shown that Yukawa couplings in a model without brane interactions are also suppressed compared with those in the standard model. It is expected that Yukawa couplings are suppressed in a more realistic model with brane interactions as well, but definitive statements must be awaited until precise form of the action is specified in the fermion sector.

Acknowledgements

This work was supported in part by JSPS fellowship No. 0509241 (Y.S.), and by Scientific Grants from the Ministry of Education and Science, Grant No. 17540257, Grant No. 13135215, Grant No. 18204024, and Grant No. 19034007(Y.H.). One of the authors (Y.H.) would like to thank the Aspen Center for Physics for its hospitality where a part of this work was done.

Appendix A

— $SO(5)$ generators —

The spinorial representation of the $SO(5)$ generators T^I is given by

$$T^{a_L} \equiv \frac{1}{2} \begin{pmatrix} \sigma_a & \\ & 0_2 \end{pmatrix}, \quad T^{a_R} \equiv \frac{1}{2} \begin{pmatrix} 0_2 & \\ & \sigma_a \end{pmatrix}, \quad T^{\hat{a}} \equiv \frac{i}{2\sqrt{2}} \begin{pmatrix} & \sigma_{\hat{a}} \\ -\sigma_{\hat{a}}^\dagger & \end{pmatrix}, \quad (\text{A.1})$$

where $T^{\hat{a}}$ ($\hat{a} = 1, 2, 3, 4$) and T^{a_L, a_R} ($a_L, a_R = 1, 2, 3$) are respectively the generators of $SO(5)/SO(4)$ and $SO(4) \sim SU(2)_L \times SU(2)_R$, and $\sigma_{\hat{a}} \equiv (\vec{\sigma}, -i1_2)$. They are normalized as

$$\text{tr}(T^I T^J) = \frac{1}{2} \delta^{IJ}, \quad (\text{A.2})$$

where $I, J = (a_L, a_R, \hat{a})$.

Appendix B

— Useful formulae for Bessel functions —

Here we collect useful formulae for the Bessel functions. $J_\alpha(z)$ and $Y_\alpha(z)$ denote the Bessel functions of the first and second kinds, respectively.

$$J_\alpha(z) \equiv \left(\frac{z}{2}\right)^\alpha \sum_{n=0}^{\infty} \frac{(-z^2/4)^n}{n! \Gamma(\alpha + n + 1)}, \quad (\text{B.1})$$

$$Y_\alpha(z) \equiv \begin{cases} \frac{1}{\sin \pi \alpha} \{ \cos \pi \alpha \cdot J_\alpha(z) - J_{-\alpha}(z) \} & \text{for } \alpha \neq \text{an integer,} \\ \frac{1}{\pi} \left[\frac{\partial J_\alpha(z)}{\partial \alpha} - (-1)^n \frac{\partial J_{-\alpha}(z)}{\partial \alpha} \right]_{\alpha=n} & \text{for } \alpha = n = \text{an integer.} \end{cases} \quad (\text{B}\cdot 2)$$

Their behavior for $|z| \gg 1$ is given by

$$\begin{aligned} J_\alpha(z) &\sim \sqrt{\frac{2}{\pi z}} \cos \left(z - \frac{(2\alpha+1)\pi}{4} \right), \\ Y_\alpha(z) &\sim \sqrt{\frac{2}{\pi z}} \sin \left(z - \frac{(2\alpha+1)\pi}{4} \right). \end{aligned} \quad (\text{B}\cdot 3)$$

These Bessel functions satisfy the following relations.

$$\begin{aligned} Z_{\alpha-1}(z) + Z_{\alpha+1}(z) &= \frac{2\alpha}{z} Z_\alpha(z), \\ \frac{dZ_\alpha(z)}{dz} &= \frac{\alpha}{z} Z_\alpha(z) - Z_{\alpha+1}(z) = Z_{\alpha-1}(z) - \frac{\alpha}{z} Z_\alpha(z), \\ J_\alpha(z)Y_{\alpha-1}(z) - Y_\alpha(z)J_{\alpha-1}(z) &= \frac{2}{\pi z}, \end{aligned} \quad (\text{B}\cdot 4)$$

$$\begin{aligned} \int^z dz \, z Z_\alpha(\lambda z) \tilde{Z}_\alpha(\lambda z) &= \frac{z^2}{4} \left\{ 2Z_\alpha(\lambda z) \tilde{Z}_\alpha(\lambda z) - Z_{\alpha-1}(\lambda z) \tilde{Z}_{\alpha+1}(\lambda z) \right. \\ &\quad \left. - Z_{\alpha+1}(\lambda z) \tilde{Z}_{\alpha-1}(\lambda z) \right\}, \end{aligned} \quad (\text{B}\cdot 5)$$

where $Z_\alpha(z)$, $\tilde{Z}_\alpha(z)$ are linear combinations of $J_\alpha(z)$ and $Y_\alpha(z)$.

Appendix C

— Properties of $F_{\alpha,\beta}(u, v)$ —

Using the Bessel functions, we define a function

$$F_{\alpha,\beta}(u, v) \equiv J_\alpha(u)Y_\beta(v) - Y_\alpha(u)J_\beta(v). \quad (\text{C}\cdot 1)$$

Using (B·4), this satisfies the following relations,

$$\begin{aligned} F_{\alpha,\beta}(u, v) &= -F_{\beta,\alpha}(v, u), \\ F_{\alpha,\alpha-1}(u, u) &= -F_{\alpha,\alpha+1}(u, u) = \frac{2}{\pi u}, \\ F_{\alpha+1,\beta}(u, v) + F_{\alpha-1,\beta}(u, v) &= \frac{2\alpha}{u} F_{\alpha,\beta}(u, v), \\ F_{\alpha-1,\alpha}(u, v) F_{\alpha,\alpha-1}(u, v) &= F_{\alpha-1,\alpha-1}(u, v) F_{\alpha,\alpha}(u, v) - \frac{4}{\pi^2 uv}. \end{aligned} \quad (\text{C}\cdot 2)$$

For non-integer α , we can express $F_{\alpha,\alpha}(u, v)$, $F_{\alpha,\alpha-1}(u, v)$, and $F_{\alpha-1,\alpha}(u, v)$ solely in terms of the Bessel function of the first kind as

$$F_{\alpha,\alpha}(u, v) = \frac{1}{\sin \pi \alpha} \{ J_{-\alpha}(u)J_\alpha(v) - J_\alpha(u)J_{-\alpha}(v) \},$$

$$\begin{aligned}
F_{\alpha,\alpha-1}(u,v) &= \frac{1}{\sin \pi \alpha} \{J_\alpha(u)J_{1-\alpha}(v) + J_{-\alpha}(u)J_{\alpha-1}(v)\}, \\
F_{\alpha-1,\alpha}(u,v) &= -\frac{1}{\sin \pi \alpha} \{J_{\alpha-1}(u)J_{-\alpha}(v) + J_{1-\alpha}(u)J_\alpha(v)\}.
\end{aligned} \tag{C.3}$$

Using these expressions and (B.1), we obtain for $z \geq 1$ and $\lambda z \ll 1$,

$$\begin{aligned}
F_{\alpha,\alpha}(\lambda, \lambda z) &= \frac{z^\alpha - z^{-\alpha}}{\pi \alpha} \{1 + \mathcal{O}(\lambda^2 z^2)\}, \\
F_{\alpha-1,\alpha}(\lambda, \lambda z) &= -\frac{2}{\pi \lambda z^\alpha} - \frac{\lambda}{2\pi \alpha(1-\alpha)} \{z^\alpha - (1-\alpha)z^{-\alpha} - \alpha z^{2-\alpha}\} \{1 + \mathcal{O}(\lambda^2 z^2)\} \\
&= -\frac{2}{\pi \lambda z^\alpha} \{1 + \mathcal{O}(\lambda^2 z^2, \lambda^2 z^{2\alpha})\}, \\
F_{\alpha,\alpha-1}(\lambda, \lambda z) &= \frac{2}{\pi \lambda z^{1-\alpha}} \\
&\quad + \frac{\lambda}{2\pi \alpha(1-\alpha)} \{z^{1-\alpha} - \alpha z^{\alpha-1} - (1-\alpha)z^{\alpha+1}\} \{1 + \mathcal{O}(\lambda^2 z^2)\} \\
&= \frac{2}{\pi \lambda z^{1-\alpha}} \{1 + \mathcal{O}(\lambda^2 z^2, \lambda^2 z^{2(1-\alpha)})\}.
\end{aligned} \tag{C.4}$$

From (B.5), we obtain

$$\begin{aligned}
\int_1^{z_\pi} dz \, z F_{\alpha,\beta}^2(\lambda_1 z, \lambda_2) &= \left[\frac{z^2}{2} \{F_{\alpha,\beta}^2(\lambda_1 z, \lambda_2) - F_{\alpha+1,\beta}(\lambda_1 z, \lambda_2) F_{\alpha-1,\beta}(\lambda_1 z, \lambda_2)\} \right]_1^{z_\pi} \\
&= \left[\frac{z^2}{2} \{F_{\alpha,\beta}^2(\lambda_1 z, \lambda_2) + F_{\alpha\pm 1,\beta}^2(\lambda_1 z, \lambda_2) \right. \\
&\quad \left. - \frac{2\alpha}{\lambda_1 z} F_{\alpha,\beta}(\lambda_1 z, \lambda_2) F_{\alpha\pm 1,\beta}(\lambda_1 z, \lambda_2) \} \right]_1^{z_\pi}.
\end{aligned} \tag{C.5}$$

In the second equality, we used the third equation of (C.2). Using this, we obtain the following integral formulae, which are useful for determining normalization factors of the mode functions.

$$\begin{aligned}
\int_1^{z_\pi} dz \, z F_{\alpha,\beta}^2(\lambda z, \lambda z_\pi) &= \frac{1}{2} \left\{ \frac{4}{\pi^2 \lambda^2} + \frac{2\alpha}{\lambda} F_{\alpha,\beta} F_{\alpha-1,\beta} - F_{\alpha,\beta}^2 - F_{\alpha-1,\beta}^2 \right\}, \\
\int_1^{z_\pi} dz \, z F_{\alpha-1,\beta}^2(\lambda z, \lambda z_\pi) &= \frac{1}{2} \left\{ \frac{4}{\pi^2 \lambda^2} + \frac{2(\alpha-1)}{\lambda} F_{\alpha,\beta} F_{\alpha-1,\beta} - F_{\alpha,\beta}^2 - F_{\alpha-1,\beta}^2 \right\}.
\end{aligned} \tag{C.6}$$

$F_{\alpha,\beta}$ in the right-hand sides are understood as $F_{\alpha,\beta}(\lambda, \lambda z_\pi)$.

References

- 1) Y. Sakamura and Y. Hosotani, Phys. Lett. B **645** (2007), 442, hep-ph/0607236.
- 2) D.B. Fairlie, Phys. Lett. B **82** (1979), 97; J. of Phys. **G5** (1979), L55.
- 3) N. Manton, Nucl. Phys. B **158** (1979), 141.
P. Forgacs and N. Manton, Commun. Math. Phys. **72** (1980), 15.
- 4) Y. Hosotani, Phys. Lett. B **126** (1983), 309.
- 5) Y. Hosotani, Ann. of Phys. **190** (1989), 233.

- 6) Y. Hosotani, Phys. Lett. B **129** (1984), 193; Phys. Rev. D **29** (1984), 731.
- 7) H. Hatanaka, T. Inami and C.S. Lim, Mod. Phys. Lett. **A13** (1998), 2601.
- 8) M. Kubo, C.S. Lim and H. Yamashita, Mod. Phys. Lett. **A17** (2002), 2249.
- 9) A. Pomarol and M. Quiros, Phys. Lett. B **438** (1998), 255.
- 10) I. Antoniadis, K. Benakli and M. Quiros, New. J. Phys. **3** (2001), 20.
- 11) C. Csaki, C. Grojean and H. Murayama, Phys. Rev. D **67** (2003), 085012.
C.A. Scrucca, M. Serone and L. Silvestrini, Nucl. Phys. B **669** (2003), 128.
- 12) L.J. Hall, Y. Nomura and D. Smith, Nucl. Phys. B **639** (2002), 307.
L. Hall, H. Murayama, and Y. Nomura, Nucl. Phys. B **645** (2002), 85.
G. Burdman and Y. Nomura, Nucl. Phys. B **656** (2003), 3.
C.A. Scrucca, M. Serone, L. Silvestrini and A. Wulzer, J. High Energy Phys. **0402** (2004), 49.
G. Panico and M. Serone, J. High Energy Phys. **0505** (2005), 024.
- 13) N. Haba, Y. Hosotani, Y. Kawamura and T. Yamashita, Phys. Rev. D **70** (2004), 015010.
N. Haba, K. Takenaga, and T. Yamashita, Phys. Lett. B **605** (2005), 355.
- 14) K.w. Choi, N. Haba, K.S. Jeong, K. Okumura, Y. Shimizu, and M. Yamaguchi, J. High Energy Phys. **0402** (2004), 037.
N. Haba and T. Yamashita, J. High Energy Phys. **0404** (2004), 016.
- 15) K. Hasegawa, C.S. Lim and N. Maru, Phys. Lett. B **604** (2004), 133.
- 16) N. Haba, K. Takenaga, and T. Yamashita, Phys. Lett. B **615** (2005), 247.
- 17) Y. Hosotani, S. Noda and K. Takenaga, Phys. Lett. B **607** (2005), 276.
- 18) G. Martinelli, M. Salvatori, C.A. Scrucca and L. Silvestrini, J. High Energy Phys. **0510** (2005), 037.
- 19) N. Maru and K. Takenaga, Phys. Rev. D **72** (2005), 046003; Phys. Lett. B **637** (2006), 287; Phys. Rev. D **74** (2006), 015017.
- 20) I. Gogoladze, T. Li, Y. Mimura and S. Nandi, Phys. Rev. D **72** (2005), 055006.
I. Gogoladze, C.A. Lee, Y. Mimura and Q. Shafi, hep-ph/0703107.
- 21) G. Cacciapaglia, C. Csaki and S.C. Park, J. High Energy Phys. **0603** (2006), 099.
- 22) G. Panico, M. Serone and A. Wulzer, Nucl. Phys. B **739** (2006), 186.
- 23) B. Grzadkowski and J. Wudka, Phys. Rev. Lett. **97** (2006), 211602.
- 24) N. Haba, S. Matsumoto, N. Okada and T. Yamashita, J. High Energy Phys. **0602** (2006), 073.
- 25) G.A. Palma, J. Phys. Conf. Ser **53** (2006), 621.
- 26) A. Aranda and J.L. Diaz-Cruz, Phys. Lett. B **633** (2006), 591.
- 27) M. Trapletti, Mod. Phys. Lett. **21** (2006), 2251.
- 28) M. Salvatori, hep-ph/0611309.
- 29) C.S. Lim, N. Maru and K. Hasegawa, hep-th/0605180.
C.S. Lim and N. Maru, hep-ph/0703017.
- 30) M. Sakamoto and K. Takenaga, Phys. Rev. D **75** (2007), 045015.
- 31) R. Contino, Y. Nomura and A. Pomarol, Nucl. Phys. B **671** (2003), 148.
- 32) K. Agashe, A. Delgado, M.J. May and R. Sundrum, J. High Energy Phys. **0308** (2003), 050.
S.J. Huber, Nucl. Phys. B **666** (2003), 269.
- 33) K. Agashe, R. Contino and A. Pomarol, Nucl. Phys. B **719** (2005), 165.
- 34) K. Agashe, G. Perez and A. Soni, Phys. Rev. D **71** (2005), 016002.
K. Agashe and R. Contino, Nucl. Phys. B **742** (2006), 59.
- 35) K. Oda and A. Weiler, Phys. Lett. B **606** (2005), 408.
- 36) Y. Hosotani and M. Mabe, Phys. Lett. B **615** (2005), 257.
- 37) Y. Hosotani, S. Noda, Y. Sakamura and S. Shimasaki, Phys. Rev. D **73** (2006), 096006.
- 38) Y. Hosotani, hep-ph/0609035.
- 39) T. Gherghetta, hep-ph/0601213.
- 40) M. Carena, E. Ponton, J. Santiago and C.E.M. Wagner, Nucl. Phys. B **759** (2006), 202; hep-ph/0701055.
- 41) A. Falkowski, Phys. Rev. D **75** (2007), 025017.
- 42) R. Contino, T. Kramer, M. Son and R. Sundrum, hep-ph/0612180.
- 43) G.F. Giudice, C. Grojean, A. Pomarol and R. Rattazzi, hep-ph/0703164.
- 44) G.v. Gersdorff, N. Irges and M. Quiros, Nucl. Phys. B **635** (2002), 127.
- 45) Y. Hosotani, in the Proceedings of “Dynamical Symmetry Breaking”, ed. M. Harada and

- K. Yamawaki (Nagoya University, 2004), p. 17. (hep-ph/0504272).
- 46) N. Irges and F. Knechtli, Nucl. Phys. B **719** (2005), 121; hep-lat/0604006; hep-lat/0609045.
 - 47) N. Maru and T. Yamashita, Nucl. Phys. B **754** (2006), 127.
 - 48) Y. Hosotani, hep-ph/0607064.
 - 49) L. Randall and R. Sundrum, Phys. Rev. Lett. **83** (1999), 3370.
 - 50) T. Gherghetta and A. Pomarol, Nucl. Phys. B **586** (2000), 141.
 - 51) S. Chang, J. Hisano, H. Nakano, N. Okada and M. Yamaguchi, Phys. Rev. D **62** (2000), 084025.
 - 52) A. Flachi, I.G. Moss, and D.J. Toms, Phys. Rev. D **64** (2001), 105029.
 - 53) C. Csaki, C. Grojean, H. Murayama, L. Pilo, and J. Terning, Phys. Rev. D **69** (2004), 055006.
 - 54) N. Sakai and N. Uekusa, hep-th/0604121.
 - 55) R.S. Chivukula, D.A. Dicus, and H.-J. He, Phys. Lett. B **525** (2002), 175.
 - 56) N. Haba, M. Harada, Y. Hosotani and Y. Kawamura, Nucl. Phys. B **657** (2003), 169 [Errata; **669** (2003) 381].
 - 57) Y. Hosotani, in the Proceedings of “*Strong Coupling Gauge Theories and Effective Field Theories*”, (Nagoya, 2002), p. 234. (hep-ph/0303066).